

# ЖУРНАЛ

гепато-гастроэнтерологических  
исследований



№3 (Том 4)

2023

# ЖУРНАЛ ГЕПАТО-ГАСТРОЭНТЕРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ

ТОМ 4, НОМЕР 3

JOURNAL OF HEPATO-GASTROENTEROLOGY RESEARCH

VOLUME 4, ISSUE 3





ISSN 2181-1008 (Online)

Научно-практический журнал  
Издается с 2020 года  
Выходит 1 раз в квартал

**Учредитель**

Самаркандский государственный  
медицинский университет,  
tadqiqot.uz

**Главный редактор:**

Н.М. Шавази д.м.н., профессор.

**Заместитель главного редактора:**

М.Р. Рустамов д.м.н., профессор.

**Ответственный секретарь**

Л.М. Гарифулина к.м.н., доцент

**Редакционная коллегия:**

Д.И. Ахмедова, д.м.н., проф;  
А.С. Бабажанов, к.м.н., доц;  
Ш.Х. Зиядуллаев, д.м.н., доц;  
Ф.И. Иноятова, д.м.н., проф;  
М.Т. Рустамова, д.м.н., проф;  
Н.А. Ярмухамедова, к.м.н., доц.

**Редакционный совет:**

Р.Б. Абдуллаев (Ургенч)  
М.Дж. Ахмедова (Ташкент)  
А.Н. Арипов (Ташкент)  
М.Ш. Ахророва (Самарканд)  
Н.В. Болотова (Саратов)  
Н.Н. Володин (Москва)  
С.С. Давлатов (Бухара)  
А.С. Калмыкова (Ставрополь)  
А.Т. Комилова (Ташкент)  
М.В. Лим (Самарканд)  
М.М. Матлюбов (Самарканд)  
Э.И. Мусабоев (Ташкент)  
А.Г. Румянцев (Москва)  
Н.А. Тураева (Самарканд)  
Ф.Г. Ульмасов (Самарканд)  
А. Фейзиоглу (Стамбул)  
Ш.М. Уралов (Самарканд)  
А.М. Шамсиев (Самарканд)  
У.А. Шербеков (Самарканд)

Журнал зарегистрирован в Узбекском агентстве по печати и информации

Адрес редакции: 140100, Узбекистан, г. Самарканд, ул. А. Темура 18.  
Тел.: +998662333034, +998915497971  
E-mail: [hepato\\_gastroenterology@mail.ru](mailto:hepato_gastroenterology@mail.ru).

## СОДЕРЖАНИЕ | CONTENT

<b>1. Атаева Ф.Н., Туразода М. У.</b> УСОВЕРШЕНСТВОВАНИЕ МЕТОДОВ ТЕРАПИИ У ДЕВОЧЕК С ЮВЕНИЛЬНЫМ КРОВОТЕЧЕНИЕМ ПРИ ЭНДОМЕТРИОЗЕ.....	5
<b>2. Garifulina L.M.,Kholmuradova Z. E. ,Kudratova G. N.</b> FEATURES OF IMPLEMENTATION OF CARDIOVASCULAR SYSTEM PATHOLOGY IN CHILDREN WITH OBESITY, IMPROVEMENT OF PREVENTION AND TREATMENT.....	8
<b>3. Гарифулина Л.М.,Тураева Д.Х.,Аслидинов Ш. Б.</b> СЕМИЗЛИГИ БЎЛГАН БОЛАЛАРДА ГЕПАТОБИЛИАР ПАТОЛОГИЯ.....	11
<b>4. Ganiev A.G., Sanakulov A.B.</b> ANALYSIS OF THE QUALITY OF LIFE OF CHILDREN WITH A PREDISPOSITION TO ALLERGIC DISEASES...	15
<b>5. Ganiev A.G., Sanakulov A.B.</b> INDICATORS OF CIRCUITAL HEART RHYTHM IN SCHOOLCHILDREN WITH NEUROCULATORY CARDIAC DYSTONIA.....	19
<b>6. Isanova Sh.T., Niyozov Sh. T., Mukhtarova M. A., Shukurov Xodixon Baxtiyor o`g`li.</b> PATHOGENETIC MECHANISM OF PAIN SYNDROME IN NEWBORNS.....	23
<b>7. Ishkabalova G.Dj.,Raxmonkulov Sh.I.</b> SURUNKALI IKKILAMCHI PIELONEFRITNI DAVOLASH XUSUSIYATLARI.....	26
<b>8. Каримжанов И.А., Мадаминова М.Ш., Умаров Д.А.</b> РОЛЬ ИНТЕРЛЕЙКИНА ИЛ-17А ПРИ ЮВЕНИЛЬНОМ ИДИОПАТИЧЕСКОМ АРТРИТЕ.....	30
<b>9. Ниязов Ш.Т., Эргашев С. С., Исанова Ш.Т., Мухтарова А. А.</b> ОПТИКО-КОГЕРЕНТНАЯ ДИАГНОСТИКА С ПЕРИНАТАЛЬНЫМИ ПАТОЛОГИЯМИ У ДЕТЕЙ.....	34
<b>10. Rasulov A. S.</b> BIOKIMYOVIY KORSATKICHLARNI O'RGANISH BOLALARDA RAXITNI DAVOLASHDA MUVAFFAQIYATGA ERISHISH YO'LIDIR.....	38
<b>11. Sirojiddinova H. N., Usmonova M. F.</b> YOSH BOLALARDA MEKONIAL ASPIRASYON SINDROMINING KLINIK XARAKTERISTIKASI.....	42
<b>12. Умаркулов З. З., Хамидов О.А., Давлатов С. С., Усмонов А.У.</b> РОЛЬ ДИАПЕВТИЧЕСКИХ МЕТОДОВ В ХИРУРГИЧЕСКОМ ЛЕЧЕНИИ КИСТОЗНЫХ ОБРАЗОВАНИЙ ПЕЧЕНИ.....	46
<b>13. Умаркулов З.З., Хамидов О.А., Давлатов С. С., Усмонов А. У.</b> РОЛЬ ДИАПЕВТИЧЕСКИХ МЕТОДОВ В ДИАГНОСТИКИ И РЕЗУЛЬТАТОВ ХИРУРГИЧЕСКОГО ЛЕЧЕНИЯ БОЛЬНЫХ ПАРАЗИТАРНЫМИ И НЕПАРАЗИТАРНЫМИ КИСТАМИ ПЕЧЕНИ.....	51
<b>14. Usmanova M.F., Sirojiddinova X. N.</b> YANGI TUG'ILGAN SHAQALOQLARDA MOSLASHUV JARAYONINING ANAMIYATI.....	56
<b>15. Xusainova Sh.K., Zakirova B.I., Makhmujanova S. R.</b> THE PREVALENCE OF RECURRENCE OF OBSTRUCTIVE BRONCHITIS IN CHILDREN.....	59
<b>16. Шавази Н.М., Ибрагимова М. Ф., Шавкатова З. Ш.</b> СОСТОЯНИЕ ЦИТОКИНОВОГО ПРОФИЛЯ У ЧАСТО БОЛЕЮЩИХ ДЕТЕЙ ПРИ ОБСТРУКТИВНОМ БРОНХИТЕ.....	63
<b>17. Шеркулов К.У., Ахророва Л.Б., Усмонова Н. У.</b> СОЧЕТАННАЯ НЕОПУХОЛЕВАЯ ПАТОЛОГИЯ ПРЯМОЙ КИШКИ И АНАЛЬНОГО КАНАЛА.....	67
<b>18. Lim M.V., Irshodzoda A. D., Xatamova K. V.</b> THE ROLE OF MICROCLIMATE IN CHILDREN WITH RECURRENT OBSTRUCTIVE BRONCHITIS.....	73
<b>19. Лим М. В., Джураева М.С., Абдурахимова А.Ф.</b> ЭФФЕКТИВНОСТЬ НЕБУЛАЙЗЕРНОЙ ТЕРАПИИ ГИАЛУРОНОВОЙ КИСЛОТОЙ У ДЕТЕЙ С БРОНХООБСТРУКТИВНЫМ СИНДРОМОМ.....	77

# JOURNAL OF HEPATO-GASTROENTEROLOGY RESEARCH

## ЖУРНАЛ ГЕПАТО-ГАСТРОЭНТЕРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ

УДК: 616.12-008 (476.2)

**Ganiev Abdurashid Ganievich**


Head of the Department of Propaedeutics  
of Childhood Diseases and Outpatient Pediatrics,  
Candidate of Medical Sciences, Associate Professor  
Andijan State Medical Institute, Andijan, Uzbekistan

**Sanakulov Abdulatif Burkhanovich**

Assistant at the Department of Childhood Diseases, Faculty of Medicine,  
Samarkand State Medical University, Samarkand, Uzbekistan

### INDICATORS OF CIRCUITAL HEART RHYTHM IN SCHOOLCHILDREN WITH NEUROCVLATORY CARDIAC DYSTONIA

**For citation:** Ganiev A.G., Sanakulov A.B./Indicators of circuit heart rhythm in school children with neurocvlatory cardiac dystonia. Journal of hepato-gastroenterology research. 2023. vol. 4, issue 3. pp.19-22

 <http://dx.doi.org/10.5281/zenodo.10511093>

#### ANNOTATION

According to the International Systematization of Diseases (10th edition), neurocirculatory dystonia (NCD) is understood as somatoform autonomic dysfunction caused by a violation of the nervous regulation of the circulatory system. Among non-infectious diseases of childhood and adolescence, the most common pathology is vegetative-vascular dystonia - from 4.8 to 29.1%. The article evaluates circadian rhythms and heart rate variability in urban and urban adolescents with similar neurocirculatory dystonia. When examining adolescents with neurocirculatory dystonia, it is necessary to conduct not only a simple ECG at rest, but also Holter ECG monitoring with determination of heart rate variability; the examination ensures adequate treatment and monitoring of patients in this group. The observed changes in circadian rhythm and rhythm variability in urban adolescents compared to rural adolescents are likely associated with living in an industrial area, environmental criteria, nutrition and greater information load, which urgently requires further research.

**Key words:** schoolchildren, neurocirculatory dystonia, cardiac rhythm and conduction disorders.

**Ganiyev Abdurashid Ganiyevich**

Bolalar kasalliklari propedevtikasi va poliklinik  
pediatriya kafedrasini mudiri, tibbiyot fanlari nomzodi, dotsent  
Andijon davlat tibbiyot instituti, Andijon, O'zbekiston

**Sanaqulov Abdulatif Burkhanovich**

Davolash fakulteti, bolalar kasalliklari kafedrasini assistenti  
Samarqand davlat tibbiyot universiteti, Samarqand, O'zbekiston

### NEYROTSIRKULYATORNI KARDIAL TURI DISTONIYASI BOR MAKTAB BOLLARIDA 24 SOATLI YURAK RITM KO'RSATKICHLARI

#### ИЗОХ.

Kasalliklarni xalqaro tizimlashtirishga ko'ra (10-nashr), neyrotsirkulyator distoni (NSD) qon aylanish tizimining asabiy regulyatsiyasi buzilishi bilan yuzaga keladigan somatoform avtonom disfunktsiyani anglatadi. Bolalik va o'smirlar davrining yuqumli bo'lmagan kasalliklari orasida vegetativ-qon tomir distoniyasi 4,8 dan 29,1% gacha bo'lgan, eng keng tarqalgan patologiya hisoblanadi. Ushbu maqolada shahar va shaharlarda yashovchi shunga o'xshash neyrokirkulyator distoni bo'lgan o'smirlarda sirkadiyalik ritm va yurak urish tezligining o'zgaruvchanligi baholanadi. Neyrotsirkulyator distoniyasi bilan og'rigan o'smirlarni tekshirishda nafaqat dam olishda oddiy EKG, balki yurak urish tezligining o'zgaruvchanligini aniqlash bilan Xolter EKG monitoringini ham o'tkazish kerak; tekshiruv ushbu guruh bemorlarini yetarli darajada davolash va monitoringini ta'minlaydi. Shahar o'smirlarida qishloq o'smirlariga nisbatan sirkadiyalik ritm va ritm o'zgaruvchanligida aniqlangan o'zgarishlar, ehtimol, sanoat zonasida yashash, atrof-muhit mezonlari, ovqatlanish va yanada ko'proq ma'lumot yuki bilan bog'liq bo'lib, bu shoshilinch ravishda keyingi tadqiqotlarni talab qiladi.

**Kalit so'zlar:** maktab o'quvchilari, neyrokirkulyator distoni, yurak ritmi va o'tkazuvchanligining buzilishi

**Ганиев Абдурашид Ганиевич**

Заведующий кафедрой пропедевтики детских болезней и поликлинической  
педиатрии, кандидат медицинских наук, доцент  
Андижанский государственный медицинский институт, Андижан, Узбекистан

**Санакулов Абдулатиф Бурханович**

**ПОКАЗАТЕЛИ КРУГОВОГО РИТМА СЕРДЦА У ШКОЛЬНИКОВ С НЕЙРОЦИРКУЛЯТОРНОЙ СЕРДЕЧНОЙ ДИСТОНИЕЙ****АННОТАЦИЯ**

Согласно Международной систематизации болезней (10-е издание) под нейроциркуляторной дистонией (НЦД) понимают соматоформную вегетативную дисфункцию, обусловленную нарушением нервной регуляции системы кровообращения. Среди неинфекционных заболеваний детского и подросткового возраста наиболее распространенной патологией является вегето-сосудистая дистония - от 4,8 до 29,1%. В статье оценены циркадные ритмы и вариабельность сердечного ритма у городских и сельских подростков со сходной нейроциркуляторной дистонией. При обследовании подростков с нейроциркуляторной дистонией необходимо проводить не только простую ЭКГ в покое, но и холтеровское мониторирование ЭКГ с определением вариабельности сердечного ритма; обследование обеспечивает адекватное лечение и наблюдение за пациентами этой группы. Наблюдаемые изменения циркадного ритма и вариабельности ритма у городских подростков по сравнению с сельскими подростками, вероятно, связаны с проживанием в промышленной зоне, экологическими критериями, питанием и большей информационной нагрузкой, что срочно требует дальнейших исследований.

**Ключевые слова:** школьники, нейроциркуляторная дистония, нарушения ритма и проводимости сердца.

According to the international systematization of diseases (10th revision), neurocirculatory dystonia (NCD) refers to somatoform autonomic dysfunction that occurs with a violation of the nervous regulation of the circulatory system [5, 6]. Among non-infectious diseases of childhood and adolescence, vegetative-vascular dystonia is considered the most common pathology with a frequency of 4.8 to 29.1% [1, 3]. It is known that vegetative-vascular dystonia or its variant neurocirculatory dystonia is considered a predictor of arterial hypertension, atherosclerosis and coronary heart disease at a young age [1, 4, 5].

The development of autonomic dysfunction in the pubertal period is facilitated by the inherent incompleteness of the morphological and active formation of the autonomic nervous system (ANS) and hormonal changes [5, 6, 7]. Neurocirculatory dystonia is considered a multifactorial disease, in the development of which genetic and acquired factors play a role, including climatic-geographical, socio-economic and household factors. It occurs in various clinical signs and syndromes, occurring in most cases under stress [7, 9, 10, 11].

The role of the autonomic nervous system in the occurrence of many types of arrhythmias, including life-threatening ones, has been convincingly confirmed in experiments with stimulation of the posterior lobe of the hypothalamus, stellate ganglia, reproduction of psychological stress, and stimulation of intracardiac parasympathetic nerves, and others.

The implementation device is associated with the influence of efferent vagal and sympathetic fibers on the electrophysiological qualities of the myocardium and the conduction system of the heart [8, 9, 11]. Currently, the majority of creators have come to a single view, namely, that one of the main pathophysiological mechanisms for the development of arrhythmias in adolescence is considered to be non-compliance with the neurogenic regulation of heart rate, leading to embodied electronic instability and myocardial instability.

In real time, determination of heart rate variability (HRV) is recognized as a more informative non-invasive way to quantify the autonomic regulation of heart rate [2, 7, 10, 11]. It is, in fact, that a decrease in HRV characteristics indicates non-compliance with the autonomic control of cardiac work, and also about the electronic instability of the myocardium. The highest HRV characteristics are recorded in healthy young people, athletes, interim - in people with various organic heart diseases, including ventricular arrhythmias, low - in people who have suffered episodes of ventricular fibrillation. The use of HRV analysis as a way to assess the adaptive probabilities of the body or the current value of stress provides practical attention for various areas of applied physiology, occupational and sports medicine.

The development of prenosological diagnostics has made it possible to distinguish among literally healthy people wide groups of people with the highest and rather high tension of regulatory systems, with an increased risk of failure of addiction and the occurrence of pathological abnormalities and diseases.

**The purpose** of this study was a comparative assessment of the daily rhythm and heart rate variability in adolescents with similar neurocirculatory dystonia, living in urban and rural areas.

**Materials and methods.** We examined 77 adolescents aged from 12 to 17 years (average age  $14.5 \pm 0.35$  years, boys -31, girls -46), who were hospitalized with a diagnosis of somatoform autonomic dysfunction, cardiac-type NCD, with January 2022 to February 2023. The information was classified according to age and area of residence (city, rural area). Taking into account the mechanisms of development of the autonomic nervous system, 2 age groups were focused. 1st age group - children 12-14 years old  $49.5 \pm 4.99\%$  (average age  $13.4 \pm 0.24$ ), of which  $62.5 \pm 8.83\%$  lived in Andijan (group IA),  $37.5 \pm 11.74\%$  of adolescents - in rural areas of Andijan region (group IB). 2nd age group - teenagers 15-17 years old  $50.5 \pm 4.99\%$  (average age  $16.5 \pm 0.29$ ), in which  $61.2 \pm 8.89\%$  of young people lived in Andijan (IIA group),  $38.8 \pm 11.49\%$  - in rural areas (IIB group). All teenagers live in the Andijan region every day.

The diagnosis of NCD was based on the aspects proposed by [6], which can be divided into personal signs and impartial data. The instrumental study included: electrocardiography (ECG) using a generally accepted method, echocardiography (ECHO-CG) with color Doppler mapping on the "Vivid 3" installation, Holter ECG monitoring with determination of heart rate variability with registration of 3 altered leads.

During daytime ECG monitoring, the characteristics of daytime dispersion of heart rate (HR) and heart rhythm disturbances were taken into account. The circadian index (CI) was calculated as the ratio of the average daytime to average nighttime heart rate (normal 1.24-1.44 a.u., average  $1.32 \pm 0.08$ ).

Heart rate variability was perceived according to the characteristics of time analysis, including these indicators, such as SDANN (standard anomaly from the average value of R-R intervals in another 5-minute fragments), SDNN (standard anomaly from the arithmetic mean), RMSSD (square root of the average sum squared differences between other R-R intervals). The frequency test was carried out on the following spectra: 1) ultra-low frequencies, ULF up to 0.003 Hz; 2) rather low frequencies, VLF -0.003-0.04 Hz; 3) low frequencies, LF 0.04-0.15 Hz; 4) the highest high frequencies, HF -0.15-0.4 Hz. The ULF and VLF spectra depend, in a key way, on the sympathetic system, HF - on the parasympathetic system, and others - on two systems at the same time. The ratio of LF to HF was interpreted as an indicator of dynamic equilibrium between two independent systems.

**Results.** Among the examined adolescents with cardiac NCD, complaints of pain in the heart area were similar (in age group I - in  $70.8 \pm 6.56\%$ , in age group II - in  $83.7 \pm 5.28\%$  of those examined). The pain, in general, had a temporary (lasting a few seconds), stabbing character. Interruptions in heart function were noticed by  $43.8 \pm 7.16\%$  of young people in group I and  $55.1 \pm 7.11\%$  of young people in age group II.

In addition,  $31.3 \pm 6.69\%$  of adolescents in age group I and  $36.7 \pm 6.89\%$  of young people in group II complained of headache, dizziness, impotence, and excessive fatigue.

In all individuals included in the study, the contractility and thickness of the walls of the ventricles of the heart, the dimensions of the ventricles and atria, the structure and function of the valve apparatus

were determined using echocardiographic techniques. More often (50.0±7.22% of those examined in age group I and 63.3±6.89% of young people in age group II), echocardiographic examination revealed slight dilatation of the heart cavities (leading to an increase in the end-diastolic volume of the left ventricle by an average of 4.1±0.5 mm).

12 (25.0±6.25%) adolescents of the 1st age group were diagnosed with mitral valve prolapse (MVP) of the 1st degree without

regurgitation, a similar diagnosis was made to 11 schoolchildren (22.5±5.97%) of the 2nd age group. Abnormal left ventricular trabeculae were observed in 29 (60.2 ± 7.07%) subjects of group I and in 30 (60.2 ± 6.99%) subjects of group II. In other adolescents, no pathology was detected during echocardiographic examination.

**Table 1**  
**Frequency of occurrence of cardiac arrhythmias and conduction disturbances in the study groups according to a normal ECG at rest (in%)**

Index	Groups			
	IA	IIA	IB	IIB
Bradycardia	26,7±8,08	43,3± 9,05	16,7±9,05	15,8±8 ,62
Tachycardia	33,3±8,6	11,1±7,62	33,3±8,61	15,8±8,62
Load on the left ventricle	33,3±8,6	5,6±5,58	30,0±8,37	2 1,1±9,62

As you can see from Table 1, in adolescents of the first age group living in a metropolis, sinus tachycardia and symptoms of stress on the left ventricle were recorded more often on a normal ECG. However, in

age group II, young people living in a metropolis were diagnosed with bradycardia more often than young people living in rural areas.

table 2

**Frequency of occurrence of heart rhythm and conduction disturbances in the study groups according to Holter ECG monitoring (in%)**

Index	Groups			
	IA	IIA	IB	IIB
Tendency to tachycardia throughout the day	63,3±8,79	22,2±10,08	36,7±8,79	15,8±8,58
Autonomic dysfunction of the sinus node	36,7±8,79	61,1±11,82	43,3±9,05	15,8±8,58

Holter monitoring data (Table 2) showed that the predisposition to sinus tachycardia in the direction of day and night in age group I was more often observed in adolescents living in a metropolis (average heart rate > 90 beats/min). Among urban young people 15-17 years old, autonomic dysfunction of the sinus node was detected more often per day. The concept of dysfunction of the sinus node includes these electrocardiographic configurations, such as sinus bradycardia (during the day and night, the average heart rate is below the age norm), pieces of movement of the pacemaker, slipping decreases, atrial rhythm throughout the day and night, slowing down of atrioventricular conduction to incomplete atrioventricular block I degree both during the day and at night, for example. In 10.2±4.32% of young people of the second age group, atrioventricular dissociation and incomplete atrioventricular block of the second degree with Samoilov-Wenckebach times were registered per day.

The circadian index in urban and rural adolescents of the first age group was on average 1.33±0.09, in the second age group - 1.38±0.08, which actually fits into the norm. The maximum average rhythm pause in children of age group I was 1.5±0.05 seconds, which actually corresponds to the upper limit of generally accepted standards; in age group II it was 1.7±0.05 seconds, i.e. older than generally accepted standards. Supraventricular extrasystole was recorded in 66.7±6.8% of urban adolescents aged 12-14 years, in rural adolescents - in 44.4±12.05% of cases. In age group II, supraventricular extrasystole was detected in 20.0±7.3% of urban young people and in 15.8±8.58% of rural ones. The largest number of extrasystoles per day is 6331, the smallest is 1. In adolescents with frequent supraventricular extrasystoles, couplets, massive extrasystoles, allorhythmia (bigeminy, trigeminy), as well as polytopic and interpolated extrasystoles were recorded. Ventricular extrasystole was observed in 26.7±8.08% of urban children aged 12-14 years and 16.7±9.05% of rural children, with the largest number per day-1429, the smallest-1. In age group II (15-17 years), ventricular extrasystole was recorded in 26.7±8.08% of urban young people and 15.8±8.59 rural ones. The maximum ventricular ectopic energy per day is -7609, the lowest is -1.

In 45.8±7.19% of children in the first age group, a clear circadian pattern of ectopic energy was noted (as in supraventricular, for example, ventricular extrasystoles), mainly in the daytime (i.e., sympathodependent ectopia), in 25.0± 6 .25% extrasystole was

recorded, for the most part, at night (vago-dependent ectopia), in others - both during the day, for example, at night. In the older age group, a reverse desire was noted: 51.0±7.14% of young people had a vagal-dependent arrhythmia, 20.4±5.76 had a sympathodependent arrhythmia, and others had a mixed type of ectopia. These configurations can be associated with the steps in the development of the function of the autonomic nervous system.

According to the SDANN and SDNN indicators, the scatter function was considered (with a sinus tempo they reflect the energy of the parasympathetic nervous system), and RMSSD is responsible for the concentration function (with a sinus tempo they reflect the energy of the sympathetic nervous system) systems).

Confinement In adolescents 12-14 years old with neurocirculatory dystonia of the cardiac type, living in a metropolis, sinus tachycardia and symptoms of stress on the left ventricle were detected more often on a normal ECG of calmness compared with adolescents of the same age living in rural areas. In the group of young people aged 15-17 years living in a metropolis, bradycardia and arrhythmia were recorded more often than in rural areas. When conducting Holter ECG monitoring in adolescents 12-14 years old living in a metropolis, a predisposition to tachycardia during the day and night was determined more often than in rural children, while in age group II, in urban young people, sinus dysfunction was diagnosed more often. node.

In adolescents aged 12-14 years with neurocirculatory dystonia of the cardiac type, living in a metropolis, compared with rural adolescents, the time SDANN indicator is truly reduced, which actually indicates a decrease in vagal energy. When assessing the spectral characteristics of the HRV, it was revealed that in urban adolescents aged 12-14 years, the indicator of sympathetic-parasympathetic tone is truly higher than generally accepted standards, which indicates the dominance of the sympathetic division of the ANS. In the older age group, a similar indicator among urban young people was truly lower than generally accepted standards, which actually indicates the dominant influence of the vagus in these adolescents.

The discovered changes in heart rate variability reflect the circadian dynamics of ectopia: the majority of adolescents 12-14 years old had sympathodependent arrhythmia, and young people 15-17 years old had vagal-dependent arrhythmia.

When examining adolescents with neurocirculatory dystonia, it is necessary to conduct not only a normal ECG at rest, but also Holter ECG monitoring with determination of heart rate variability; the examination will ensure adequate treatment and supervision of the given group of patients.

The discovered changes in daily rhythm and rhythm variability in urban adolescents compared with rural ones are probably associated with living in an industrial zone, with the criteria of the environment, nutrition, and an even greater information load, which urgently requires further research.

#### Список литературы/ Iqtiboslar / References

1. Aleksandrov, A. A. Prevention of cardiovascular diseases from youth: plans, success, problems / A. A. Aleksandrov // *Cardiology*. 2005.No. 7. P. 4-8.
2. Heart rate variability test when using electrocardiographic systems (methodological recommendations) / R. M. Baevsky [et al.] // *Bulletin of Arrhythmology*. 2011.No. 24. S. 66-85.
3. Belokon, N. A. Diseases of the heart and blood vessels in children / N. A. Belokon, M. B. Kuberge. M.: Medicine, 2007. pp. 136 -197.
2. Belokon, N.A. Vegetative-vascular dystonia in children: hospital, diagnosis, treatment / N. A. Belokon, G. G. Osokina, I. V. Leontyeva.M. : Medicine, 2007.P. 24.
3. Vein, A. M. Autonomic disorders: hospital, healing, diagnosis / A. M. Vein. M.: Medical Information Agency, 2008. P. 49-87.
4. 6.Ganiev A.G., Zainabitdinova S.N. Daily blood pressure monitoring is the main way to diagnose arterial hypertension in adolescents//re-health journal No. 2. (14)-2022 art.: 148-153
5. Ganiev A.G. Mitral valve prolapse in children with connective tissue dysplasia // *Quarterly scientific and practical journal // "Hepato-gastroenterological studies"* No. 4. What 3. 2022. Art. 14-16.
6. Азимова К. Т., Гарифулина Л. М. Показатели цитокинового статуса при бронхиолите у детей // *Journal of cardiorespiratory research*. – 2023. – Т. 1. – №. 2. – С. 45-47.6.
7. Гойибова Н. С., Гарифулина Л. М. semizligi bor bolalarda buyraklar va ularning funksional holati // *Журнал гепато-гастроэнтерологических исследований*. – 2022. – Т. 3. – №. 1.
8. Жамшедовна А. М., Гарифулина Л. М. Samarqand viloyatidagi semizligi bor bolalarda vitamin d statusini baholash // *Журнал гепато-гастроэнтерологических исследований*. – 2023. – Т. 4. – №. 2.



# ЖУРНАЛ ГЕПАТО-ГАСТРОЭНТЕРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ

ТОМ 4, НОМЕР 3

JOURNAL OF HEPATO-GASTROENTEROLOGY RESEARCH

VOLUME 4, ISSUE 3

**Editorial staff of the journals of [www.tadqiqot.uz](http://www.tadqiqot.uz)**

Tadqiqot LLC The city of Tashkent,  
Amir Temur Street pr.1, House 2.

Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Phone: (+998-94) 404-0000

**Контакт редакций журналов. [www.tadqiqot.uz](http://www.tadqiqot.uz)**

ООО Tadqiqot город Ташкент,  
улица Амира Темура пр.1, дом-2.

Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Тел: (+998-94) 404-0000