

**CRR**  
JOURNAL  
OF CARDIORESPIRATORY RESEARCH

**ISSN 2181-0974**  
**DOI 10.26739/2181-0974**



**Journal of**  
**CARDIORESPIRATORY**  
**RESEARCH**



Volume 2, Issue 2

**2021**

МИНИСТЕРСТВО ЗДРАВООХРАНЕНИЯ  
РЕСПУБЛИКИ УЗБЕКИСТАН

Журнал кардиореспираторных исследований

# JOURNAL OF CARDIORESPIRATORY RESEARCH

Главный редактор: Э.Н.ТАШКЕНБАЕВА

Учредитель:

Самаркандский государственный  
медицинский институт

Tadqiqot.uz

Ежеквартальный  
научно-практический  
журнал



ISSN: 2181-0974

DOI: 10.26739/2181-0974



**N° 2  
2021**

**Главный редактор:**

**Ташкенбаева Элеонора Негматовна**

*доктор медицинских наук, заведующая кафедрой внутренних болезней №2 Самаркандского Государственного Медицинского института, председатель Ассоциации терапевтов Самаркандской области. <https://orcid.org/0000-0001-5705-4972>*

**Заместитель главного редактора:**

**Хайбулина Зарина Руслановна**

*доктор медицинских наук, руководитель отдела биохимии с группой микробиологии ГУ «РСНПМЦХ им. акад. В. Вахидова» <https://orcid.org/0000-0002-9942-2910>*

**ЧЛЕНЫ РЕДАКЦИОННОЙ КОЛЛЕГИИ:**

**Аляви Анис Лютфуллаевич**

*академик АН РУз, доктор медицинских наук, профессор, Председатель Ассоциации Терапевтов Узбекистана (Ташкент) <https://orcid.org/0000-0002-0933-4993>*

**Бокерия Лео Антонович**

*академик РАН и РАМН, доктор медицинских наук, профессор, Президент научного центра сердечно-сосудистой хирургии им. А.Н. Бакулева (Москва), <https://orcid.org/0000-0002-6180-2619>*

**Курбанов Равшанбек Давлатович**

*академик АН РУз, доктор медицинских наук, профессор, Председатель Республиканского специализированного научно-практического медицинского центра кардиологии (Ташкент) <https://orcid.org/0000-0001-7309-2071>*

**Michał Tendera**

*профессор кафедры кардиологии Верхнесилезского кардиологического центра, Силезский медицинский университет в Катовице, Польша (Польша) <https://orcid.org/0000-0002-0812-6113>*

**Покушалов Евгений Анатольевич**

*доктор медицинских наук, профессор, заместитель генерального директора по науке и развитию сети клиник «Центр новых медицинских технологий» (ЦНМТ), (Новосибирск), <https://orcid.org/0000-0002-2560-5167>*

**Акилов Хабибулла Атауллаевич**

*доктор медицинских наук, профессор, ректор Ташкентского Института Усовершенствования Врачей (Ташкент)*

**Абдиева Гулнора Алиевна**

*ассистент кафедры внутренних болезней №2 Самаркандского Медицинского Института <https://orcid.org/0000-0002-6980-6278> (ответственный секретарь)*

**Ризаев Жасур Алимджанович**

*доктор медицинских наук, профессор, Ректор Самаркандского государственного медицинского института <https://orcid.org/0000-0001-5468-9403>*

**Зиядуллаев Шухрат Худойбердиевич**

*доктор медицинских наук, доцент, проректор по научной работе и инновациям Самаркандского Государственного медицинского института <https://orcid.org/0000-0002-9309-3933>*

**Зуфаров Миржамол Мирумарович**

*доктор медицинских наук, профессор, руководитель отдела ГУ «РСНПМЦХ им. акад. В. Вахидова» <https://orcid.org/0000-0003-4822-3193>*

**Ливерко Ирина Владимировна**

*доктор медицинских наук, профессор, заместитель директора по науке Республиканского специализированного научно-практического медицинского центра фтизиатрии и пульмонологии Республики Узбекистан (Ташкент) <https://orcid.org/0000-0003-0059-9183>*

**Цурко Владимир Викторович**

*доктор медицинских наук, профессор кафедры общей врачебной практики Института профессионального образования Первого Московского государственного медицинского университета им. И.М. Сеченова (Семёновский университет) (Москва) <https://orcid.org/0000-0001-8040-3704>*

**Тураев Феруз Фатхуллаевич**

*доктор медицинских наук, главный научный сотрудник отделения приобретенных пороков сердца ГУ «РСНПМЦХ им. акад. В. Вахидова»*

**Насирова Зарина Акбаровна**

*PhD, ассистент кафедры внутренних болезней №2 Самаркандского Государственного Медицинского Института (ответственный секретарь)*

**Bosh muharrir:**

**Tashkenbayeva Eleonora Negmatovna**

*tibbiyot fanlari doktori, Samarqand davlat tibbiyot instituti 2-sonli ichki kasalliklar kafedrasini mudiri, Samarqand viloyati vrachlar uyushmasi raisi.*  
<https://orsid.org/0000-0001-5705-4972>

**Bosh muharrir o'rinbosarlari:**

**Xaibulina Zarina Ruslanovna**

*tibbiyot fanlari doktori, "akad V. Vohidov nomidagi RIJM davlat institutining mikrobiologiya guruhi bilan biokimyoviy kafedrasini mudiri"*  
<https://orcid.org/0000-0002-9942-2910>

**TAHRIRIYAT A'ZOLARI:**

**Alyavi Anis Lyutfullayevich**

*O'zbekiston Respublikasi Fanlar akademiyasining akademigi, tibbiyot fanlari doktori, professor, O'zbekiston Terapevtlar uyushmasi raisi (Toshkent),*  
<https://orcid.org/0000-0002-0933-4993>

**Bockeria Leo Antonovich**

*Rossiya fanlar akademiyasining akademigi, tibbiyot fanlari doktori, professor, A.N. Bakuleva nomidagi yurak-qon tomir jarrohligi ilmiy markazi prezidenti (Moskva)*  
<https://orcid.org/0000-0002-6180-2619>

**Qurbanov Ravshanbek Davlatovich**

*O'zbekiston Respublikasi Fanlar akademiyasining akademigi, tibbiyot fanlari doktori, professor, Respublika ixtisoslashtirilgan kardiologiya ilmiy-amaliy tibbiyot markazining raisi (Toshkent)*  
<https://orcid.org/0000-0001-7309-2071>

**Mixal Tendera**

*Katovitsadagi Sileziya Tibbiyot Universiteti, Yuqori Sileziya Kardiologiya Markazi kardiologiya kafedrasini professori (Polsha)*  
<https://orcid.org/0000-0002-0812-6113>

**Pokushalov Evgeniy Anatolevich**

*tibbiyot fanlari doktori, professor, "Yangi tibbiy texnologiyalar markazi" (YTTM) klinik tarmog'ining ilmiy ishlar va rivojlanish bo'yicha bosh direktorining o'rinbosari (Novosibirsk)*  
<https://orcid.org/0000-0002-2560-5167>

**Akilov Xabibulla Ataulayevich**

*tibbiyot fanlari doktori, professor, Toshkent vrachlar malakasini oshirish instituti rektori (Toshkent)*

**Abdiyeva Gulnora Aliyevna**

*Samarqand davlat tibbiyot instituti 2-sonli ichki kasalliklar kafedrasini assistenti (mas'ul kotib)*  
<https://orcid.org/0000-0002-6980-6278>

**Rizayev Jasur Alimjanovich**

*tibbiyot fanlari doktori, professor, Samarqand davlat tibbiyot instituti rektori*  
<https://orcid.org/0000-0001-5468-9403>

**Ziyadullayev Shuxrat Xudoyberdiyevich**

*tibbiyot fanlari doktori, dotsent, Samarqand davlat tibbiyot institutining fan va innovatsiyalar bo'yicha prorektori (Samarqand)*  
<https://orcid.org/0000-0002-9309-3933>

**Zufarov Mirjamol Mirumarovich**

*tibbiyot fanlari doktori, professor, "akad V. Vohidov nomidagi RIJM davlat muassasasi" bo'limi boshlig'i "*  
<https://orcid.org/0000-0003-4822-3193>

**Liverko Irina Vladimirovna**

*tibbiyot fanlari doktori, professor, Respublika ixtisoslashtirilgan ftiziologiya va pulmonologiya ilmiy-amaliy tibbiyot markazining ilmiy ishlar bo'yicha direktor o'rinbosari (Toshkent)*  
<https://orcid.org/0000-0003-0059-9183>

**Surko Vladimir Viktorovich**

*I.M. Sechenov nomidagi Birinchi Moskva Davlat Tibbiyot Universiteti Kasbiy ta'lim institutining umumiy amaliyot shifokorlik amaliyoti kafedrasini tibbiyot fanlar doktori, professori (Semyonov universiteti) (Moskva)*  
<https://orcid.org/0000-0001-8040-3704>

**To'rayev Feruz Fatxullayevich**

*tibbiyot fanlari doktori, "akad V. Vohidov nomidagi RIJM davlat institutining yurak kasalliklari" bo'limining bosh ilmiy xodimi*  
<https://orcid.org/0000-0002-1321-4732>

**Nosirova Zarina Akbarovna**

*Samarqand davlat tibbiyot instituti 2-sonli ichki kasalliklar kafedrasini assistenti, PhD (mas'ul kotib)*

**Chief Editor:**

**Tashkenbaeva Eleonora Negmatovna**

*Doctor of Medical Sciences, Head of the Department of Internal Diseases No. 2 of the Samarkand State Medical Institute, Chairman of the Association of Physicians of the Samarkand Region. <https://orsid.org/0000-0001-5705-4972>*

**Deputy Chief Editor:**

**Xaibulina Zarina Ruslanovna**

*Doctor of Medical Sciences, Head of the Department of Biochemistry with the Microbiology Group of the State Institution "RSSC named after acad. V. Vakhidov", <https://orcid.org/0000-0002-9942-2910>*

**MEMBERS OF THE EDITORIAL BOARD:**

**Alyavi Anis Lutfullaevich**

*Academician of the Academy of Sciences of the Republic of Uzbekistan, Doctor of Medical Sciences, Professor, Chairman of the Association of Physicians of Uzbekistan (Tashkent) <https://orcid.org/0000-0002-0933-4993>*

**Bockeria Leo Antonovich**

*Academician of the Russian Academy of Sciences, Doctor of Medical Sciences, Professor, President of the Scientific Center for Cardiovascular Surgery named after A.N. Bakuleva (Moscow) <https://orcid.org/0000-0002-6180-2619>*

**Kurbanov Ravshanbek Davlatovich**

*Academician of the Academy of Sciences of the Republic of Uzbekistan, Doctor of Medical Sciences, Professor, Chairman of the Republican Specialized Scientific and Practical Medical Center of Cardiology, (Tashkent) <https://orcid.org/0000-0001-7309-2071>*

**Michal Tendera**

*Professor of the Department of Cardiology, Upper Silesian Cardiology Center, Silesian Medical University in Katowice, Poland (Poland) <https://orcid.org/0000-0002-0812-6113>*

**Pokushalov Evgeny Anatolyevich**

*Doctor of Medical Sciences, Professor, Deputy Director General for Science and Development of the Clinic Network "Center for New Medical Technologies" (CNMT), (Novosibirsk) <https://orcid.org/0000-0002-2560-5167>*

**Akilov Xabibulla Ataulaevich**

*Doctor of Medical Sciences, Professor, Rector of the Tashkent Institute for the Improvement of Physicians (Tashkent)*

**Abdieva Gulnora Alieva**

*Assistant of the Department of Internal Diseases No. 2 of the Samarkand State Medical Institute <https://orcid.org/0000-0002-6980-6278>  
(Executive Secretary)*

**Rizaev Jasur Alimjanovich**

*Doctor of Medical Sciences, Professor, Rector of the Samarkand State Medical Institute <https://orcid.org/0000-0001-5468-9403>*

**Ziyadullaev Shuhrat Khudoyberdievich**

*Doctor of Medical Sciences, Associate Professor, Vice-Rector for Science and Innovation of the Samarkand State Medical Institute (Samarkand) <https://orcid.org/0000-0002-9309-3933>*

**Zufarov Mirjamol Mirumarovich**

*Doctor of Medical Sciences, Professor, Head of the Department of the State Institution "RSNPMTSH named after acad. V. Vakhidov" <https://orcid.org/0000-0003-4822-3193>*

**Liverko Irina Vladimirovna**

*Doctor of Medical Sciences, Professor, Deputy Director for Science of the Republican Specialized Scientific and Practical Medical Center for Phthisiology and Pulmonology of the Republic of Uzbekistan (Tashkent) <https://orcid.org/0000-0003-0059-9183>*

**Tsurko Vladimir Viktorovich**

*Doctor of Medical Sciences, professor of the Department of General Medical Practice of First Moscow State Medical University by name I.M. Sechenov (Sechenov University) (Moscow) <https://orcid.org/0000-0001-8040-3704>*

**Turaev Feruz Fatxullaevich**

*Doctor of Medical Sciences, Chief Researcher of the Department of Acquired Heart Diseases of the State Institution "RSNPMTSH named after acad. V. Vakhidov"*

**Nasirova Zarina Akbarovna**

*PhD, Assistant of the Department of Internal Diseases No. 2 of the Samarkand State Medical Institute (Executive Secretary)*

**ЧЛЕНЫ РЕДАКЦИОННОГО СОВЕТА | TAHRIRIYAT KENGASHI**  
**MEMBERS OF THE EDITORIAL BOARD:**

**Алимов Дониёр Анварович**  
доктор медицинских наук, директор  
Республиканского научного центра  
экстренной медицинской помощи

**Янгиев Бахтиёр Ахмедович**  
кандидат медицинских наук,  
директор Самаркандского филиала  
Республиканского научного центра  
экстренной медицинской помощи

**Абдуллаев Акбар Хатамович**  
доктор медицинских наук, главный  
научный сотрудник Республиканского  
специализированного научно-  
практического центра медицинской  
терапии и реабилитации  
<https://orcid.org/0000-0002-1766-4458>

**Агабабян Ирина Рубеновна**  
кандидат медицинских наук, доцент,  
заведующая кафедрой терапии ФПДО,  
Самаркандского Государственного  
медицинского института

**Алиева Нигора Рустамовна**  
доктор медицинских наук, заведующая  
кафедрой Госпитальной педиатрии №1  
с основами нетрадиционной медицины  
ТашПМИ

**Исмаилова Адолат Абдурахимовна**  
доктор медицинских наук, профессор,  
заведующая лабораторией  
фундаментальной иммунологии  
Института иммунологии геномики  
человека АН РУз

**Камалов Зайнитдин Сайфутдинович**  
доктор медицинских наук, профессор,  
заведующий лабораторией  
иммунорегуляции Института  
иммунологии и геномики  
человека АН РУз

**Каюмов Улугбек Каримович**  
доктор медицинских наук, профессор,  
заведующий кафедрой внутренних  
болезней и телемедицины  
Ташкентского Института  
Усовершенствования Врачей

**Хусинова Шоира Акбаровна**  
кандидат медицинских наук, доцент,  
заведующая кафедрой общей практики,  
семейной медицины ФПДО  
Самаркандского Государственного  
медицинского института

**Alimov Doniyor Anvarovich**  
tibbiyot fanlari doktori, Respublika  
shoshilinch tibbiy yordam ilmiy  
markazi direktori (Toshkent)

**Yangiyev Baxtiyor Axmedovich**  
tibbiyot fanlari nomzodi,  
Respublika shoshilinch tibbiy  
yordam ilmiy markazining  
Samarqand filiali direktori

**Abdullaev Akbar Xatamovich**  
tibbiyot fanlari doktori, O'zbekiston  
Respublikasi Sog'liqni saqlash  
vazirligining "Respublika  
ixtisoslashtirilgan terapiya va tibbiy  
reabilitatsiya ilmiy-amaliy  
tibbiyot markazi" davlat  
muassasasi bosh ilmiy xodimi  
<https://orcid.org/0000-0002-1766-4458>

**Agababayan Irina Rubenovna**  
tibbiyot fanlari nomzodi, dotsent,  
DKTF, terapiya kafedrasini mudiri,  
Samarqand davlat tibbiyot instituti

**Alieva Nigora Rustamovna**  
tibbiyot fanlari doktori, 1-sonli  
gospital pediatriya kafedrasini mudiri,  
ToshPTI

**Ismoilova Adolat Abduraximovna**  
tibbiyot fanlari doktori, professor,  
O'zbekiston Respublikasi Fanlar  
akademiyasining Odam genomikasi  
immunologiyasi institutining  
fundamental immunologiya  
laboratoriyasining mudiri

**Kamalov Zaynitdin Sayfutdinovich**  
tibbiyot fanlari doktori, professor,  
O'zbekiston Respublikasi Fanlar  
akademiyasining Immunologiya va  
inson genomikasi institutining  
Immunogenetika laboratoriyasi mudiri

**Qayumov Ulug'bek Karimovich**  
tibbiyot fanlari doktori, professor,  
Toshkent vrachlar malakasini oshirish  
institutining ichki kasalliklar va  
teletibbiyot kafedrasini mudiri

**Xusinova Shoira Akbarovna**  
tibbiyot fanlari nomzodi, dotsent,  
Samarqand davlat tibbiyot instituti  
DKTF Umumiy amaliyot va oilaviy  
tibbiyot kafedrasini mudiri (Samarqand)

**Alimov Doniyor Anvarovich**  
Doctor of Medical Sciences, Director of  
the Republican Scientific Center of  
Emergency Medical Care

**Yangiev Bakhtiyor Axmedovich**  
PhD, Director of Samarkand branch of  
the Republican Scientific Center of  
Emergency Medical Care

**Abdullaev Akbar Xatamovich**  
Doctor of Medical Sciences, Chief  
Researcher of the State Institution  
"Republican Specialized Scientific and  
Practical Medical Center for Therapy and  
Medical Rehabilitation" of the Ministry of  
Health of the Republic of Uzbekistan,  
<https://orcid.org/0000-0002-1766-4458>

**Agababayan Irina Rubenovna**  
PhD, Associate Professor, Head of the  
Department of Therapy, FAGE,  
Samarkand State Medical Institute

**Alieva Nigora Rustamovna**  
Doctor of Medical Sciences, Head of the  
Department of Hospital Pediatrics No. 1  
with the basics of alternative  
medicine, TashPMI

**Ismoilova Adolat Abduraximovna**  
doctor of Medical Sciences, Professor,  
Head of the Laboratory of Fundamental  
Immunology of the Institute of  
Immunology of Human Genomics of the  
Academy of Sciences of the  
Republic of Uzbekistan

**Kamalov Zaynitdin Sayfutdinovich**  
Doctor of Medical Sciences, Professor,  
Head of the Laboratory of  
Immunogenetics of the Institute of  
Immunology and Human Genomics of the  
Academy of Sciences of the  
Republic of Uzbekistan

**Kayumov Ulugbek Karimovich** Doctor  
of Medical Sciences, Professor, Head of  
the Department of Internal Diseases and  
Telemedicine of the Tashkent Institute for  
the Advancement of Physicians

**Khusinova Shoira Akbarovna**  
PhD, Associate Professor, Head of the  
Department of General Practice, Family  
Medicine FAGE of the  
Samarkand State Medical Institute

Page Maker | Верстка | Sahifalovchi: Xurshid Mirzahmedov

Контакт редакций журналов. [www.tadqiqot.uz](http://www.tadqiqot.uz)  
ООО Tadqiqot город Ташкент,  
улица Амира Темура пр.1, дом-2.  
Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Телефон: +998 (94) 404-0000

Editorial staff of the journals of [www.tadqiqot.uz](http://www.tadqiqot.uz)  
Tadqiqot LLC the city of Tashkent,  
Amir Temur Street pr.1, House 2.  
Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Phone: (+998-94) 404-0000

# MUNDARIJA | СОДЕРЖАНИЕ | CONTENT

## ОБЗОРНЫЕ СТАТЬИ/ ADABIYOTLAR SHARHI/ REVIEW ARTICLES

- 1 **Alyavi B.A., Abdullaev A.H., Dalimova D.A., Uzokov J.K.**  
Питание - важный фактор профилактики и лечения сердечно-сосудистых заболеваний  
Diet as an important factor in the prevention and treatment of cardiovascular diseases.  
Yurak-qon tomir kasalliklarining oldini olishda va davolashda ratsional ovqatlanish muhim omil sifatida..... 9
- 2 **Ellamonov S.N., Tashkenbaeva E.N., Abdieva G.A., Nasyrova Z.A., Khamidov N.S.**  
Факторы прогрессирования артериальной гипертензии у больных в коморбидности с сахарным диабетом 2 типа.  
Factors of arterial hypertension progression in patients in comorbidity with type 2 diabetes mellitus.  
Qandli diabet 2 turi bilan birga hamroh kasalliklari bor bemorlarda arterial gipertenziyaning avj olish omillari..... 16
- 3 **Mullabaeva G.U., Jumaniyazov D.K.**  
Значение нового биомаркера st2 в развитии сердечной недостаточности.  
Significance of the new st2 biomarker in the development of heart failure.  
Yangi st2 biomarkerining yurak etishmovchiligini rivojlanishidagi ahamiyati..... 22
- 4 **Sachin Kumar B., Tashkenbaeva E.N., Abdieva G.A.**  
Влияние курения на сердечно-сосудистые функции: роль никотина и монооксида углерода  
Effects of smoking on cardiovascular function: the role of nicotine and carbon monoxide.  
Chekishning yurak-qon tomir funktsiyasiga ta'siri: nikotin va uglerod monoksidining roli..... 26
- 5 **Tashkenbaeva N.F., Alimova D.A., Trigulova R.Kh., Abdullaev T.A., Alimov A.V.**  
Взаимосвязи сердечной недостаточности и сахарного диабета.  
Interrelation of heart failure and diabetes mellitus.  
Yurak etishmovchiligi va qandli diabetning o'zaro bogliqligi..... 34
- 6 **Tashkenbaeva E.N., Nasyrova Z.A., Saidov.M.A.,**  
Механизмы кардиопротективных эффектов десфлурана и севофлурана во время реперфузии.  
Stratification of chronic ischemic heart disease depending on diagnostic methods and ways of their treatment.  
Desfluran va sevofluranning kardioprotektiv ta'sirlari mexanizmalari..... 39
- 7 **Teshaev Sh.J., Dzhumaev K.Sh., Razhabova K.Sh.**  
Влияние вредных привычек и физической активности на группах пожилых и старческих возрастах.  
Effect of hazardous habits and physical activity on senior and senior ages.  
Zararli odatlar va jismoniy faollikning keksa va qari yoshli aholi guruhiga ta'siri..... 43
- 8 **Togaeva B.M., Tashkenbaeva E.N., Abdieva G.A., Khaydarova D.D. Abdullaeva Z.A.**  
Течение covid-19 у больных с кардиоваскулярными заболеваниями.  
COVID-19 course in patients with cardiovascular diseases.  
Covid-19 yurak qon tomir kasalliklari bor bemorlarda kechishi..... 47

## ORIGINAL MAQOLALAR/ОРИГИНАЛЬНЫЕ СТАТЬИ/ORIGINAL ARTICLES

- 9 **Alavi A.L., Kenjayev S.R., Nazarova M.X., Kenjaev S.R., Kaxarov I.I., Latipov N.M.**  
Оценка систолической функции левого желудочка и клинических показателей в зависимости результатов реперфузии у больных инфарктом миокарда с подъемом ST.  
Assessment of systolic function and indicators of left ventricular remodeling depending on the results of reperfusion in patients with ST elevation myocardial infarction.  
ST-elevatsiya bilan infarkti bo'lgan bemorlarda reperfuzyiya natijalariga qarab chap qorincha sistolik funktsiyasini va klinik parametrlarini baholash..... 51
- 10 **Kadirova G.G., Raimkulova N.R., Mamajanova Z.Sh., Ikramova A.Sh., Ernazarova M.M.**  
Структурно-гемодинамические особенности сердца при артериальной гипертензии у женщин в период менопаузы во взаимосвязи с психоэмоциональным статусом.  
Structural and hemodynamic features of the heart in arterial hypertension in women in period of menopause in interconnection with psych-emotional status.  
Arterial gipertenziya bilan hastalangan menopauza davridagi ayollarda psixoemotsional status bilan birgalikdagi yurakning struktur-gemodinamik xususiyatlari..... 58
- 11 **Karimova N.A., Kurbanova N.S.**  
Нарушение физического развития у подростков и его осложнения.  
Developmental impairment in adolescents and its complications.  
O'spirinlarda rivojlanayotgan buzilish va uning asoratlari..... 64

12	<b>Salivonchik D., Salivonchik E.</b> Энергетический коллапс при covid-19: новые маркеры диагностики, аспекты лечения. Energy COVID-19 collapse: new diagnostic markers, treatment aspects. COVID-19 da energya kollapsi: diagnostikaning yangi markerlari, davolanish aspektlari.....	68
13	<b>Tulaboeva G.M., Talipova Yu.Sh., Abdukodirova N.M., Kamolov B.B., Saidov Kh.Kh.</b> Клинико-гемодинамическая эффективность холина альфосцерата при артериальной гипертензии и гипертонической энцефалопатией. Clinical and hemodynamic efficacy of choline alfoscerate in arterial hypertension and hypertensive encephalopathy. Arterial gipertoniya va gipertonik entsefalopatiyada xolin alfostseratning klinik gemodinamik samaradorligi.....	77
14	<b>Khodzhaeva S.A., Adzhablaeva D.N.</b> Факторы риска и качество жизни у детей с латентной туберкулезной инфекцией. Risk factors and life`s quality condition of children with latent tuberculous infection. Latent sil infeksiyasi bor bolalarda havf omillari hamda hayot sifatining holati.....	82
15	<b>Khasanjanova F.O., Tashkenbaeva E.N., Boltakulova S.D.</b> Факторы риска, влияющие на течение нестабильных вариантах стенокардии у мужчин в молодом и пожилом возрасте с дислипидемией. Risk factors affecting the course of unstable angina in young and elderly men with dyslipidemia. Dislipidemiya bo'lgan yosh va keksa yoshdagi erkaklarda stenokardiyaning beqaror variantlarini kechishida xavf omillarining ta'siri.....	87
16	<b>Xoljigitova M.B., Safarova M.P., Niyozova F.N., Okboev T.A., Rustamova Sh.Sh., Jurayev S.O.</b> Кардиоваскулярные изменения сердечно-сосудистой системы у больных с хронической обструктивной болезнью легких. Cardiovascular changes in the cardiovascular system in patients with chronic obstructive pulmonary disease. Surunkali obstruktiv o'pka kasalligi bo'lgan bemorlarda yurak-qon tomir tizimidagi yurak-qon tomir o'zgarishlar.....	92
17	<b>Yakubbekov N.T.</b> Агрегация тромбоцитов у больных с многососудистым поражением коронарного русла на фоне сахарного диабета. Plate aggregation in patients with multivesel coronary artery diseases loss in the background of diabetes mellitus. Qandli diabet fonida ko'p toj tomirlar shikastlanishiga ega bemorlarda trombositlar agregatsiasi.....	99





# JOURNAL OF CARDIORESPIRATORY RESEARCH

## ЖУРНАЛ КАРДИОРЕСПИРАТОРНЫХ ИССЛЕДОВАНИЙ

**Сачин Кумар Байга**

Студент MBBS

Самаркандский Государственный Медицинский Институт  
Самарканд, Узбекистан

**Ташкенбаева Элеонора Негматовна**


д.м.н., проф., зав. кафедрой внутренних болезней №2  
Самаркандский государственный медицинский институт  
Самарканд, Узбекистан

**Абдиева Гулнора Алиевна**

ассистент кафедры внутренних болезней №2  
Самаркандский государственный медицинский институт  
Самарканд, Узбекистан

### ВЛИЯНИЕ КУРЕНИЯ НА СЕРДЕЧНО-СОСУДИСТЫЕ ФУНКЦИИ: РОЛЬ НИКОТИНА И МОНООКСИДА УГЛЕРОДА

**For citation:** Sachin Kumar B., Tashkenbaeva E.N., Abdieva G.A. Effects of smoking on cardiovascular function: the role of nicotine and carbon monoxide. Journal of cardiorespiratory research. 2021, vol.2, issue 2, pp.26-33

 <http://dx.doi.org/10.26739/2181-0974-2021-2-4>

### АННОТАЦИЯ

Курение — это распространенное явление среди всех факторов сердечно-сосудистых заболеваний. Он влияет на миокард и его кровоснабжение, усиливает атеросклероз и способствует развитию инфаркта миокарда, кардиомиопатии и заболеваний периферических сосудов. Никотин, окись углерода и различные элементы табака оказывают непосредственное воздействие на эндотелий, вызывая воспаление, атерому и тромбоз. В сигаретном дыме много свободных радикалов, которые провоцируют и усиливают воспалительный каскад, увеличивая инфильтрацию лейкоцитов и производство цитокинов. Очень важно собирать правильные записи о привычке к курению, однако накопление записей больше не должно зависеть от собственных записей, которые вводят в заблуждение, однако следует использовать биохимические биомаркеры нескольких форм, в идеале котинин. Курильщики должны получить интенсивные консультации, каждый с помощью квалифицированного персонала на месте и с помощью профессиональных служб по отказу от курения. Курильщикам следует дать фундаментальное определение пагубным последствиям курения для ишемической болезни сердца и атеросклероза. Пациентам следует рекомендовать применить все повторные рекомендации по консультированию и обратиться к заместительной терапии или другим лекарственным средствам, чтобы бросить курить. Цель этой статьи - дать краткую оценку результатов курения, и особенно результатов воздействия никотина и других веществ на сердечно-сосудистую систему. Никотин деактивирует сердечную вегетативную функцию, усиливает эмпатию, увеличивает частоту сердечных сокращений (ЧСС) при расслаблении, в то же время замедляя повышение частоты сердечных сокращений за счет инновационных упражнений и снижает максимальное количество часов, которое может быть выполнено. В то же время образующийся курением ко связывается с гемоглобином и миоглобином, снижает сатурацию артериальной крови O<sub>2</sub>, ставит под угрозу работу респираторных ферментов и вызывает нарушение устройства производства, транспортировки и транспортировки O<sub>2</sub>, особенно при выполнении упражнений, в частности, снижая целевые возможности и общая производительность циркуляционной машины.

**Ключевые слова:** болезнь сердца, сердечно-сосудистая система, курение, табак.

**Sachin Kumar Baitha**

MBBS Student

Samarkand State Medical Institute  
Samarkand, Uzbekistan

**Tashkenbaeva Eleonora Negmatovna**

Doctor of Medical Sciences, Prof., Head, Department  
of Internal Medicine №2

Samarkand State Medical Institute  
Samarkand, Uzbekistan

**Abdieva Gulnora Alieva**

Assistant of the Department of Internal Medicine No. 2  
Samarkand State Medical Institute  
Samarkand, Uzbekistan

## EFFECTS OF SMOKING ON CARDIOVASCULAR FUNCTION: THE ROLE OF NICOTINE AND CARBON MONOXIDE

## ANNOTATION

Smoking is a widespread chance thing for all factors of cardiovascular disorder. It influences the myocardium and occludes the blood supply, will increase atherosclerosis and contributes to myocardial infarction, cardiomyopathy and peripheral vascular disorder. Nicotine, carbon monoxide and different tobacco elements have direct outcomes at the endothelium, inflicting inflammation, atheroma and thrombosis. Free radicals are ample in cigarette smoke and those provoke and accentuate the inflammatory cascade, growing leukocyte infiltration and cytokine production. It is essential to reap correct records approximately smoking habit, however records amassing ought to now no longer depend on self-record that is misleading, however ought to use a few shapes of biochemical biomarker, ideally cotinine. Smokers ought to then be intensively counselled, each through in-residence skilled personnel and thru the professional smoking cessation services. It ought to be defined to sufferers in fundamental phrases approximately the deleterious outcomes of smoking on coronary heart disorder and atherosclerosis. Patients ought to be recommended to apply all reasserts of counselling and to go to the substitute treatments or different pharmaceutical aids to quit. The purpose of this article is to provide a brief evaluation of the outcomes of smoking, and especially the outcomes of nicotine and co on cardiovascular function. Nicotine deactivates cardiac autonomic function, will increase empathy hobby, increases heart rate (HR) at relaxation, while blunting HR elevation all through innovative exercise and lowering the maximum HR that may be executed. On the equal time, the smoking-generated co binds with haemoglobin and myoglobin, reduces arterial O<sub>2</sub> blood saturation, compromises the performance of respiratory enzymes, and reasons disorder of the O<sub>2</sub> production, transportation and transport device, specially all through exercise, notably decreasing the purposeful capability and the overall performance of the circulatory machine.

**Keywords:** heart disease, circulatory system, smoking, tobacco.

**Sachin Kumar Baita**

MBBS yo'nalishi bo'yicha talaba  
Samarqand davlat tibbiyot instituti  
Samarqand, O'zbekiston

**Tashkenbayeva Eleonora Negmatovna**

Tibbiyot fanlari doktori, prof., Ichki kasalliklar  
kafedrasini mudiri №2  
Samarqand davlat tibbiyot instituti  
Samarqand, O'zbekiston

**Abdiyeva Gulnora Aliyevna**

2-son ichki kasalliklar kafedrasini assistenti  
Samarqand davlat tibbiyot instituti  
Samarqand, O'zbekiston

## CHEKISHNING YURAK-QON TOMIR FUNKTSIYASIGA TA'SIRI: NIKOTIN VA UGLEROD MONOKSIDINING ROLI

## ANNOTATSIIYA

Chekish yurak-qon tomir kasalliklarining barcha omillari uchun keng tarqalgan imkoniyatdir. Bu miyokardga ta'sir qiladi va qon bilan ta'minlanmaydi, aterosklerozi kuchaytiradi va miyokard infarkti, kardiomyopatiya va periferik qon tomir kasalliklariga yordam beradi. Nikotin, uglerod oksidi va turli xil tamaki elementlari endoteliyda to'g'ridan-to'g'ri natijalarga ega bo'lib, yallig'lanish, atheroma va tromboz keltirib chiqaradi. Erkin radikallar sigaretaning tutunida juda ko'p bo'lib, ular yallig'lanish kaskadini qo'zg'atadi va ta'kidlaydi, leykotsitlar infiltratsiyasi va sitokin ishlab chiqarishni ko'paytiradi. Taxminan chekishni odat qilish uchun to'g'ri yozuvlarni yig'ish kerak, ammo hozirda to'plangan yozuvlar chalg'ituvchi o'z-o'zini yozib olishga bog'liq emas, ammo bir nechta biokimyoviy biomarker, ideal holda kotinindan foydalanish kerak. Keyinchalik chekuvchilarga har birida yashash joyidagi malakali kadrlar va chekishni tashlash bo'yicha professional xizmatlar orqali intensiv ravishda maslahat berish kerak. Chalinganlarga yurakning koronar buzilishi va ateroskleroz kasalligi bo'yicha chekishning zararli natijalarini asosiy iboralar bilan aniqlash kerak. Bemorlarga maslahatning barcha qayta tiklanishlarini qo'llash va davolanishni to'xtatish uchun almashtirish muolajalariga yoki turli xil farmatsevtik vositalarga murojaat qilish tavsiya etilishi kerak. Ushbu maqolaning maqsadi chekish natijalarini qisqacha baholash, xususan yurak-qon tomir faoliyati bo'yicha nikotin va ko'pinchalarini baholashdir. Nikotin yurakning avtonom funktsiyasini o'chiradi, hamdardlik sevimli mashg'ulotlarini kuchaytiradi, bo'shashganda yurak urishini (soat) oshiradi, shu bilan birga soat ko'tarilishini innovatsion mashqlar yordamida va bajarilishi mumkin bo'lgan maksimal soatni pasaytiradi. Xuddi shu vaqtda, chekish natijasida hosil bo'lgan CO gemoglobin va miyoglobin bilan bog'lanib, qonning qon bilan to'yinganligini pasaytiradi, nafas olish fermentlarining ish faoliyatini pasaytiradi va O<sub>2</sub> ishlab chiqarish, tashish va tashish moslamasining buzilishini, xususan, jismoniy mashqlar orqali, shu bilan birga maqsadga muvofiq qobiliyat va qon aylanish mashinasining umumiy ishlashi.

**Kalit so'zlar:** yurak kasalligi, qon aylanish tizimi, chekish, tamaki

## Introduction


Smoking is a major hazard thing for cardiovascular morbidity and mortality, and is considered to be the main preventable purpose of loss of life within the global [61, 65]. Internationally, 25% of centre - aged cardiovascular deaths because of smoking [7]. The European society of cardiology said these days that smoking reasons 28% of cardiovascular deaths in men aged 35 to 69 years and 13% in girls of the identical age [40]. In the European region of the arena fitness organization (WHO), smoking is the second one most vital chance issue inside the burden of incapacity adjusted life years and is the primary danger aspect for untimely mortality, related to about 1.6 million deaths every yr. [66]. In the eu union (EU), 15 % of all -cause deaths are attributed smoking, amounting to 655,000 smoking associated deaths each yr. [53]. At the same time as in Greece, the share of deaths from any smoking related cause, among individuals a while 35 and older, has been envisioned to

be 18.1%. Primarily based on estimates via the who, tobacco maintains to kill nearly 6 million humans each year - together with more than 600,000 passive people who smoke – via coronary heart ailment, lung cancer, and different ailments [65]; that is one and a 1/2 million extra than the corresponding estimate for 1990 [7]. If contemporary tendencies continue, the death toll is projected to attain extra than 8 million according to yr. by using 2030 [65].

Smoking ranks among the pinnacle causes of cardiovascular ailment, together with coronary heart disease, ischemic stroke, peripheral artery disorder and abdominal aortic aneurysm [40]. It's also associated with an improved chance of sure styles of most cancers, and is a primary purpose of chronic obstructive pulmonary disease [61, 65]. Smoking, both lively or passive, can cause cardiovascular ailment thru a chain of interdependent approaches, consisting of superior oxidative pressure, hemodynamic and autonomic changes, disorder,

inflammation, hyperlipidemia, or different outcomes [11]. Endothelial thrombosis, even exposure to small portions-e.g., Occasional smoking, passive smoking, some cigarettes in keeping with day-is enough to have deleterious results [60]. Cigarette smoke carries extra than 4000 chemical substances which have dangerous effects on cardiovascular function [17]. These include nicotine, carbon monoxide (CO), oxidative gases, polycyclic fragrant hydrocarbons, carbonyls, butadiene, minerals, carbon disulphide, and benzene. Even though the various toxic materials contained in tobacco smoke are ordinary merchandise of the combustion of natural substances, exposure to smoking entails contact with materials which can be unique to tobacco smoke and are regarded to be unfavorable to the fitness: nicotine and co [62].

No matter all the exposure regarding the documented unfavorable consequences of smoking on public health, smoking prevalence nevertheless remains excessive in the ecu, where about 30% of citizens are people who smoke. Greece suffers from a widespread smoking-associated public health hassle, having the very best share of people who smoke within the EU (42%). In keeping with who estimates, in Greece 63.1% of grownup males and 39% of ladies  $\geq 15$  years are people who smoke [67], while in young adults, aged 19-30 years, the smoking incidence is 37,6% [51], the motive of this article is to provide a quick description of the results of smoking, and especially the outcomes of nicotine and co, on cardiovascular characteristic, supplying essential statistics that would make a contribution to decreasing the smoking epidemic and its effects for cardiovascular health.



Nicotine

Nicotine causes your blood vessels to constrict or narrow, that limits the number of blood that flows to your organs. Over time, the constant constriction leads to blood vessels that are stiff and fewer elastic. Constricted blood vessels decrease the amount of atomic number 8 associated nutrients your cells receive. to satisfy the requirement for a lot of oxygen, your pulse could increase. phytotoxin is classed as an alkaloid (like opiate and cocaine) and meets the factors of an extremely addictive drug. One coffin nail delivers 1.2-2.9 mg of nicotine, and also the typical one pack-per-day smoker absorbs 20-40 mg of nicotine on a daily basis [43]. As associate addictive drug, phytotoxin has a pair of terribly potent effects: it's a stimulant and it is additionally a depressant [28]. phytotoxin deregulates viscus function, sixteen sympathetic activations, raises heart rate, causes coronary and involuntary boosts peripheral vasoconstriction, will increase cardiac muscle workload, and stimulates adrenal and neuronic hormone release [3]. In addition, nicotine is related to hypoglycemic agent resistance, increased humor super molecule levels, and intravascular inflammation that contributes to the event of atherosclerosis [3].

#### Vascular Function

There are sample published data suggesting that prolonged exposure to tobacco smoke causes pathological alteration of endothelial function. Endothelial dysfunction can be caused by metabolic (dyslipidemia), environmental (smoking), and physiological (arterial hypertension) factors, or by inflammation that provokes pathological conditions [41]. It is characterized by an imbalance between vasodilatory and vasoconstrictive substances derived from the endothelium, anticoagulant and procoagulant mechanisms, growth factors and growth inhibitors [41]. Under normal circumstances, free radicals circulating in the human body are neutralized by defensive mechanisms. However, if their concentrations within the blood must rise greatly due to excessive exposure to harmful factors such as smoking, they cannot be controlled and dangerous mutations that destroy cells can occur. Oxidative stress is seen to arise under these conditions [17].

The term "oxidative stress" refers to the whole of the intracellular and extracellular situations that result in chemical or metabolic manufacturing of reactive oxygen species (ROS) [62]. Smoke exists specifically in states: the gaseous (which incorporates CO) and the solid

(tar). In each those states, it consists of a big number of unfastened radicals [11]. Pryor and Stone decided that 1 g of tar from cigarette smoke consists of greater than 10 long-lived unfastened radicals (hours to months), while 1 g risky fraction of smoke consists of 10 [3] short-lived unfastened radicals (seconds). Chronic publicity to tobacco additionally weakens the antioxidant shielding mechanisms that alter those big numbers of smoking-caused unfastened radicals, main to a sizeable growth in oxidative stress [41]. Oxidative stress, the oxidation of lipids, proteins, and DNA, is without delay related to atherogenesis [17]. An indicative locating is that once stages of isoprostanes (indexes of lipid peroxidation and oxidative damage) have been measured in smokers, their stages have been determined to be better than in non-smokers [62]. The response of nitric oxide (NO) with the unfastened radicals contained in smoke reduces NO's bioavailability, interfering with its vasodilatory, antithrombotic, anti-inflammatory, and antioxidant effects, in addition to its have an impact on endothelium permeability and myocardial function (lowering the diastolic distensibility of the left ventricle) [35]. The alteration in biosynthesis of NO and its reduced activity [8], in mixture with the smoking-caused discount in prostacyclin manufacturing [22] and the direct poisonous impact of nicotine on endothelial cells that reasons direct structural damage [3], are crucial elements which can result in endothelial dysfunction Using an extract of cigarette tobacco or its remoted ingredients, along with nicotine, many in vitro research have determined that smoking is related to decreased NO availability. It has been proven that nicotine attention in smokers' blood serum reduces the supply of NO in human umbilical vein endothelial cells (HUVECs), in addition to in human coronary artery endothelial cells, main to a discount withinside the brachial artery's endothelium-established vasodilation [11]. Using this model, Barua et al. confirmed that publicity to smokers' sera reduced NO availability in each HUVECs and human coronary artery endothelial cells, with the aid of using changing the expression and pastime of the endothelial NO synthase enzyme [8]. In addition, they cited a large correlation among glide-mediated brachial artery endothelium-established vasodilation and NO bioavailability from cultured HUVECs uncovered to serum from the identical individuals. On the alternative hand, CO, that is extensively expanded in smokers, inhibits the advent of NO and takes its vicinity in hemoglobin bonds [27]. These findings result in the realization that the big portions of unfastened radicals contained in smoke decorate oxidative pressure and, in aggregate with decreased NO bioavailability, nicotine-triggered vasoconstriction and impaired vasodilation, can also additionally result in endothelial dysfunction.

The consequent endothelial harm contributes to the formation and development of atheromatous plaque, and decreases blood glide thru thrombosis and vasospasm, therefore inflicting cardiovascular disease [5, 41].

#### Lipid Metabolism

Tobacco smoke, and especially nicotine, has a significant effect on lipid metabolism and the regulation of lipid levels in the blood [46]. Therefore, cigarette smoke may promote atherosclerosis partly through its effects on the lipid profile [11]. It has been shown that, in the presence of already increased serum lipid levels, smoking Event [33].

Smoking is related to extensively increased serum concentrations of general LDL cholesterol and triglycerides [46]. In addition, numerous researches have proven a bent for low-density lipoprotein (LDL) and really low-density lipoprotein (VLDL) LDL cholesterol to be barely better in smokers [47]. These institutions appear to be dose dependent [46]. On the alternative hand, smoking lowers serum concentrations of high-density lipoprotein (HDL) LDL cholesterol, an effective shielding issue in opposition to the improvement of atherosclerosis [64]. The distinction is generally small, five mg/dl or less, however this distinction represents a 10crease and could be predicted to have an effect on atherogenesis to a great degree [47]. Giving up smoking improves HDL levels, irrespective of frame weight, contributing to a development in cardiovascular fitness after smoking cessation [45].

It is feasible that oxidative harm to protein and lipid components may also give an explanation for the manner wherein cigarette smoke impacts plasma LDL and HDL. Cigarette smoking will increase the oxidative change of LDL. Exposure to cigarette smoke extract

additionally decreases the plasma activity of paraoxonase, an enzyme that protects in opposition to LDL oxidation [11]. There are capability mechanisms through which reactive smoke additives can produce their deleterious results on critical plasma components: 1) indirectly, fueloline-section cigarette smoke may also spark off macrophages and neutrophils within the lung, which may also launch enzymes and oxidants able to adverse lipids and proteins; 2) directly, for the reason that lung possesses a really big floor place for fueloline exchange, it's miles feasible that fueloline-section cigarette smoke additives have interaction with plasma components within the interstitial fluid [64].

#### Arteriosclerosis

Arteriosclerosis is a general term that includes almost all arterial disorders that cause thickening and hardening of all types of arteries. Atherosclerosis is a specific form of arteriosclerosis, the most characteristic feature of which is the concentration of lipids in the intima of large elastic arteries (aorta) and medium-sized muscular arteries (coronary, femoral, carotid, etc.) [45]. Smoking is considered a significant risk factor for the development of atherosclerosis. The atherosclerotic effects of cigarette smoke are largely due to events related to thrombosis. The accumulation of platelets coating the artery wall in sites where there is turbulent blood flow or endothelial injury may be the prodromal stage for the formation of atheromatous plaque [54].

Nicotine is idea to be chargeable for the boom in blood viscosity and platelet aggregation, because it inhibits the manufacturing of prostacyclin which might restrict platelet aggregation [33]. Increases manufacturing of platelet adhesion thrombi, divides coronary artery intima, quickens the procedure of atheromatous plaque formation, and is related to an extended threat of cardiac ischemia [5]. In addition, nicotine influences prostaglandin metabolism, weakening the vessel's defense towards platelet deposition [56]. The boom in platelet aggregation, the impact of nicotine on blood coagulation time, and the boom in blood viscosity, in aggregate with the boom in stages of LDL and VLDL, the discount in HDL, and inflammatory processes, sell the advent of atheromatous plaque and the improvement of atherosclerosis [3, 28, 47]. It is hence probable that continual smoking, via way of means of growing peripheral vascular resistances on this way, can also additionally result in a boom in cardiac afterload and a consequent discount in stroke volume [50].

The circulating degree of fibrinogen in people who smoke is one of the maximum effective predictive markers of coronary events.

The boom in fibrinogen tiers acts in mixture with the boom in purple mobileular mass from long-time period publicity to CO, growing blood viscosity and boosting the activation of platelets, for this reason growing the chance of atherogenesis. Increased fibrinogen tiers within the blood flow also can result in the improvement of atherosclerosis, with an instantaneous impact at the boom in platelet aggregation [39].

Tissue factor (TF)—in any other case referred to as tissue platelet issue, or issue III, or thrombokinase, or CD142—is a protein discovered in endothelial tissue, platelets, and leucocytes, and is important for the initiation of thrombus formation with the aid of using zymogen prothrombin [21]. TF is expressed with the aid of using cells which can be commonly now no longer uncovered to blood flow, together with sub-endothelial cells (e.g. smooth-muscle cells) and the cells that surround blood vessels (e.g. fibroblasts). This can extrade whilst blood vessels are damaged—for instance with the aid of using bodily injury, or rupture, or atherosclerotic plaque. TF is found in atherosclerotic plaque and might sell thrombogenesis and likely propagation of the thrombus to the already present atherosclerosis. Sambola et al. discovered that smoking multiplied plasma TF ranges in people who smoke who smoked 10 or greater cigarettes consistent with day, with a smoking record of 10 or more years [57].

Atherogenesis and coronary artery disorder are the end result of inflammatory processes. The reality that smoking is related to infection means that infection can be one of the mechanisms thru which cigarette smoking ends in cardiovascular dysfunction. C-reactive protein (CRP) and degrees of white blood cells are markers of infection, and are hence related to atherosclerosis and an extended chance of cardiovascular disorder [2]. Levels of CRP and white blood cells appear like better in people who smoke than in non-people who smoke [24]. Furthermore,

there seems to be a relation among the volume of smoking and the white blood mobility count. Dietrich et al. claimed that the boom in CRP found in people who smoke is proportional to each the amount and the years of smoking [24].

Overall, Nicotine increases sympathetic activity, stimulates the release of neurotransmitters, causes coronary and peripheral vasoconstriction, and elevates blood pressure.

In addition, nicotine enhances lipolysis, increases free fatty acid levels, increases oxidative stress, endothelial damage and dysfunction, and promotes vessel inflammation, contributing significantly to the development of atherosclerosis and heart disease.

#### Autonomic Nervous System

There is a long-time hyperlink among atypical coronary heart rate (HR) responses at relaxation and throughout exercise, autonomic disorder and cardiovascular health [57]. On the opposite hand, persistent smoking is related to disorder of the autonomic anxious device [10, 63], and the atypical HR responses to tobacco can be implicated within the hyperlink among smoking and cardiovascular disease [4, 20]. Although the ideal mechanism of movement of smoke components remains beneath Neath investigation, all proposed hypotheses nation that the principal outcomes of smoking on cardiovascular characteristic are related to the direct or oblique movement of nicotine at the nutrireulation of the circulatory device, in which sympathetic hobby is accelerated and parasympathetic hobby is reduced.

The nicotine-caused sympathetic overdrive reasons the adrenal medulla to boom the secretion of each epinephrine and norepinephrine into the circulating blood [4]. In addition, nicotine stimulates the vasomotor center of the medulla, inflicting secretion of norepinephrine from neighborhood deposits. Subsequently, secretion of catecholamines from the loose nerve endings of the sympathetic nerves and the neighborhood launch of epinephrine and norepinephrine are accelerated. In addition, vasoconstriction of coronary vessels occurs, the biosynthetic ability of prostacyclin is reduced, and endothelial characteristic is impaired [58]. The stimulation of catecholamine secretion, in mixture with the depressed manufacturing of prostacyclin (strong vasodilators), outcomes in an acute upward thrust in blood pressure, a substantial upward thrust in HR, an boom in cardiac contractility, and a substantial boom in myocardial work. Nicotine impacts cardiovascular characteristic each directly, as defined previously, and indirectly, via a sequence of neurohormonal changes [9]. In particular, nicotine molecules engage with and prompt the brain's acetylcholine receptors (nAChRs), whose extended activation might also additionally desensitize a percentage of them. The activation of nAChRs via way of means of nicotine boosts the discharge of neurotransmitters, whilst changing the characteristic of a number of them—which includes norepinephrine, dopamine, serotonin (5-HT), and endogenous opioid peptides—accordingly enhancing the movement of the peripheral anxious device and inflicting cigarette addiction [15].

#### Heart Rate at Rest

Smoking has been related to expanded resting HR values in wholesome adults, irrespective of age or sex. Minami et al. located that the HR is on common 7 bpm quicker in people who smoke than in non-people who smoke [13]; this locating is consistent with records mentioned via way of means of Papanthassiou et al., indicating that people who smoke had a drastically better resting HR as compared with non-people who smoke in each female (76.4 bpm vs. 70.0 bpm,  $p=0.001$ ) and male (72.8 vs. 66.3,  $p=0.004$ ) subjects [50]. In particular, smoking is related to selective changes in cardiac autonomic control [51]. More specifically, smoking, performing at peripheral sympathetic sites, will increase circulating tiers of catecholamines, augments sympathetic outflow, and reasons a long-time period discount in vagal drive. This sympathetic predominance, visible even in younger heavy people who smoke, is likewise related to impaired baroreflex function, main to a marked boom in HR at rest.

#### Heart Rate during Exercise

During exercising, the accelerated metabolic needs are met via way of means of an accelerated cardiac output, carried out via an augmentation in HR and stroke volume. The elevation of HR, that is related to age, HR relaxation and exercising capacity, is regulated via way of means of exercising-caused autonomic control, in which

sympathetic hobby will increase and vagal tone is reduced. The HR elevation peaks at maximal exercising, whilst healthful topics acquire a real HR max close ( $\pm 10$  bpm) to their age-anticipated HR max [32]. An impaired HR reaction to exercising and failure to reach  $>80\%$  of the age-anticipated HR max, referred to as chronotropic incompetence, are related to autonomic imbalance and are vital prognostic markers of cardiovascular health. In many HR-associated researches in healthful adults, smoking become located to blunt HR elevation throughout innovative exercising and to decrease the most HR carried out, posing an accelerated hazard to smokers' health. diversifications to continual smoking, inclusive of down law of  $\beta$ -adrenergic receptors, were used to provide an explanation for smokers' blunted HR reaction to exercising. long-time period smoking has been located to lower the density of lymphocyte or adipose tissue  $\beta$ -receptors, down-regulating the  $\beta$ -receptors of the cardiovascular system [29].

#### Heart rate recovery

On the final touch of full of life exercising, sympathetic pastime withdraws and vagal reactivation mediates the price at which HR decreases, making the post-exercising HR decline a crucial reference marker for cardiac autonomic control. attenuated HR decline for the duration of healing is a crucial surrogate for underlying autonomic disorder this is related to extended cardiovascular morbidity and mortality [30]. in lots of epidemiological HR-associated research in wholesome middle-elderly populations, smoking became inversely related to HR decline for the duration of healing.

#### Smoking and insulin resistance

Insulin has an impact on almost all of the tissues of the body, both immediately or indirectly, and is characterized as a garage hormone due to its anabolic motion on all 3 important nutritional groups: namely, carbohydrates, fats, and proteins [25].

Insulin is related to precise receptors within the cell membrane. the primary capabilities of the insulin receptor are to recognize and to bind the hormone with the goal cell, and to transmit its metabolic action. if any such capabilities are disturbed, insulin resistance is manifested. insulin resistance, metabolic syndrome, and glucose intolerance are seemed as disturbances with a not unusual place history and sturdy interrelations [55].

Nicotine is known to growth sympathetic activity, to elevate circulating degrees of catecholamines, increase hormone, adrenocorticotropic hormone, cortisol, prolactin, and beta-endorphin, and to lower estrogen degrees these kinds of consequences are strongly adversarial to insulin's action. Thus, smoking reduces insulin production, slowing glucose catabolism and main to its accumulation within the body [36]. Nicotine might also growth insulin resistance directly. It has been proven that the growth in insulin resistance become halted after nicotine substitute become stopped, or even progressed throughout non-stop weight benefit, implying that nicotine as opposed to weight benefit can be answerable for the preliminary growth in insulin resistance found in a few smoking-associated studies. The smoking-brought on insulin resistance is likewise related to an growth in triglyceride count, due to the fact in fatty tissue glucose is transformed to triglycerides [1]. In turn, because of multiplied serum concentrations of FFA and triglycerides, insulin-inspired glucose delivery in skeletal muscle of routine cigarette people who smoke is exceptionally impaired in contrast with non-people who smoke. Insulin resistance and the growth in triglycerides found in people who smoke are sizable threat elements for the destiny improvement of arteriosclerosis and as a result cardiovascular disease [12].

#### Carbon Monoxide

Carbon monoxide (CO) is made from the unfinished combustion of carbon-containing substances, inclusive of fuel and tobacco. The history stage of CO within the surroundings may be very low and has little impact on humans, even as maximum of the CO produced with the aid of using herbal or technological tactics is oxidized to CO<sub>2</sub> within the top surroundings. Comparatively, then, the 3-6% CO in cigarette smoke (and the 2-3 instances better concentrations in pipe and cigar smoke) constitute notably better tiers than are generally encountered [20].

Carbon monoxide publicity has been implicated within the procedure of atherosclerosis, contributing to the buildup of LDL

cholesterol within the aorta and coronary arteries. In addition, CO publicity complements endothelial damage, main to destructive outcomes within the presence of ischemic coronary heart or peripheral vascular disorder. The deleterious outcomes of CO are extra profound within the myocardium than in peripheral tissues, due to the very excessive oxygen extraction with the aid of using the myocardium at rest [26]. There is epidemiological proof that employees uncovered to excessive CO concentrations have better cardiovascular morbidity and mortality as compared to the predicted charge within the preferred population. The foremost mechanism with the aid of using which CO reasons coronary heart disorder is thru hypoxia. Inhalation of cigarette smoke, with the aid of using both lively or passive smokers, will increase the tiers of carboxyhemoglobin (COHb) within the blood, lowering the delivery of O<sub>2</sub> to the tissues. In addition, myoglobin binds with CO in order that the coronary heart muscle does now no longer take in the essential O<sub>2</sub> and does now no longer carry out optimally. The decreased O<sub>2</sub> uptake due to smoking, collectively with a growth in serum lactic acid tiers (lactic acidosis), ends in a discount in top cardio capability and to a great lower in most O<sub>2</sub> uptake (VO<sub>2</sub>max) [49].

#### CO and Haemoglobin

The robust chemical affinity among haemoglobin (Hb) and CO is well-known. It has been predicted that the affinity among Hb and CO is two hundred instances extra than the affinity among Hb and oxygen (O<sub>2</sub>). A direct outcome of this distinction is the considerable binding of Hb via way of means of CO within the blood, the advent of COHb [34], and a awesome growth in its serum levels, ensuing in a extensive lower in oxygen uptake via way of means of peripheral tissues. More specifically, the CO in smoke binds Hb, developing COHb thru the subsequent reaction:



#### CO and Myoglobin

Myoglobin can also additionally integrate with CO and, like Hb, has a more affinity (30-50 times) with CO than with O<sub>2</sub>, intensifying the hypoxaemia of peripheral tissues and mainly the lively muscles. However, myoglobin binds to 1 molecule of O<sub>2</sub>, while Hb binds to four. Thus, the bad consequences of improved COHb tiers are a lot greater hanging than the ones of COMb, efficiently lowering each the O<sub>2</sub> deliver to the tissues and the O<sub>2</sub> uptake of running muscles.

#### CO and lactic acidosis

The term "lactic acidosis" refers to excessive stages of lactic acid within the blood. The decreased performance of the O<sub>2</sub> transportation and deliver device in people who smoke inhibits mitochondrial function. The publicity of mitochondria to smoking-brought on oxidative materials consequences in harm to the mitochondrial DNA, lowering adenosine triphosphate manufacturing in coronary heart and muscle cells [66]. Essentially, smoking disturbs the enzyme activity (adenine nucleotide translocator and mitochondrial superoxide dismutase) in mitochondria this is crucial for his or her proliferation, for this reason lowering their numbers. Because of this harm, the muscular tissues can't get the electricity they want to function (on account that they not have enough mitochondria); they consequently are seeking for electricity thru some other route: anaerobic metabolism [52]. The latter process, however, has lactic acid as its very last product, so that the amount of circulating lactic acid will increase significantly (lactic acidosis), growing the blood's acidity, compromising cardio tolerance, and impairing workout capacity [6].

#### CO and Exercise Capacity

Smoking even one cigarette can right now have an effect on bodily workout ability. The results of CO, including the considerable binding of Hb and the decreased arterial O<sub>2</sub> blood saturation, the insufficiency of respiration enzymes, in aggregate with the binding of myoglobin and the results of CO on cardio metabolism, bring about disorder of the O<sub>2</sub> manufacturing, transportation, and transport system, especially for the duration of workout [19]. Briefly, the decreased portions of transported O<sub>2</sub> and the reduced O<sub>2</sub> deliver to and uptake from the energetic tissues, mixed with the binding of myoglobin via way of means of CO, appreciably lower maximal oxygen uptake (VO<sub>2</sub>max) lowering the

purposeful ability and the overall performance of the circulatory system [37].

There is an observable lower, of round 10%, within the period of workout till exhaustion in smokers, that is on account of a discount in O<sub>2</sub> manufacturing within the metabolically energetic tissues, due to arterial O<sub>2</sub> desaturation, and to the insufficiency of the O<sub>2</sub> transportation, delivery and uptake system [14]. This impaired workout tolerance and the reduced maximal workout ability were recorded even in younger wholesome smokers.

Similar outcomes of smoke at the O<sub>2</sub> transportation and delivery system are visible in people who aren't lively people who smoke. Since non-people who smoke are extra liable to CO than people who smoke, absolutely being uncovered to cigarette smoke may also lessen their VO<sub>2</sub>max. The quantity to which VO<sub>2</sub>max is decreased relies upon on the quantity of CO that people who smoke inhale. Horvath et al. claimed that no massive discount in VO<sub>2</sub>max became discovered till stages of COHb reached or surpassed 4.3%, a degree exhibited with the aid of using maximum people who smoke. 100 From the instant COHb stages attain 4.3%, VO<sub>2</sub>max decreases [23] according with the subsequent equation

$$VO_{2max} = 0.91(\%COHb) + 2.2$$

## Conclusions

Smoking, through its primary components nicotine and CO, will increase oxidative stress, endothelial harm and disorder, is related to appreciably better serum concentrations of overall ldl cholesterol and triglycerides, reduces the aerobic defensive HDL, and via way of means of selling intravascular irritation represents a big hazard issue for the improvement of atherosclerosis and cardiovascular disease [48]. In addition, nicotine deregulates cardiac autonomic function, boosts sympathetic activity, and will increase HR at rest, whilst it blunts HR elevation all through modern workout and lowers the most HR that may be achieved. In parallel, the smoking-caused CO binds with haemoglobin and myoglobin, reduces arterial O<sub>2</sub> blood saturation, and compromises the performance of respiration enzymes, ensuing in disorder of the O<sub>2</sub> production, transportation and shipping system, specifically all through workout; this will extensively lessen the purposeful potential and the overall performance of the circulatory system [38]. Altogether, smoking is the maximum essential modifiable hazard issue for cardiovascular disease, a first-rate hazard issue for cardiovascular morbidity and mortality, and is taken into consideration to be the main preventable reason of demise within the world [18].

## Список литературы/ Iqtiboslar/References

- Alves-Bezerra M, Cohen DE. Triglyceride Metabolism in the Liver. *Compr Physiol*. 2017;8(1):1-8. Published 2017 Dec 12. doi:10.1002/cphy.c170012
- Asthana A, Johnson HM, Piper ME, Fiore MC, Baker TB, Stein JH. Effects of smoking intensity and cessation on inflammatory markers in a large cohort of active smokers. *Am Heart J*. 2010; 160(3):458-463
- Atherosclerosis. Author manuscript; available in PMC 2013 Aug 28. Published in final edited form as: *Atherosclerosis*. 2011 Apr; 215(2): 281–283. Published online 2011 Feb 1. doi: 10.1016/j.atherosclerosis.2011.01.003
- Audrey A. Wickiser, Plasma catecholamine and ascorbic acid levels in smokers and nonsmokers as a function of stress, University of Nebraska at Omaha DigitalCommons@UNO, 5-1984
- Badimon L, Padró T, Vilahur G. Atherosclerosis, platelets and thrombosis in acute ischaemic heart disease. *Eur Heart J Acute Cardiovasc Care*. 2012;1(1):60-74. doi:10.1177/2048872612441582 42:1149–1160.
- Baker JS, McCormick MC, Robergs RA. Interaction among Skeletal Muscle Metabolic Energy Systems during Intense Exercise. *J Nutr Metab*. 2010;2010:905612. doi:10.1155/2010/905612.
- Banks, E., Joshy, G., Korda, R.J. et al. Tobacco smoking and risk of 36 cardiovascular disease subtypes: fatal and non-fatal outcomes in a large prospective Australian study. *BMC Med* 17, 128 (2019). <https://doi.org/10.1186/s12916-019-1351-4> (last day accessed 3 July 2019)
- Barua RS, Ambrose JA, Eales-Reynolds LJ, DeVoe MC, Zervas JG, Saha DC. Dysfunctional endothelial nitric oxide biosynthesis in healthy smokers with impaired endothelium-dependent vasodilatation. *Circulation*. 2001; 104:1905-1910.
- Benowitz NL, Burbank AD. Cardiovascular toxicity of nicotine: Implications for electronic cigarette use. *Trends Cardiovasc Med*. 2016;26(6):515-523. doi:10.1016/j.tem.2016.03.001
- Benowitz NL. Cigarette smoking and cardiovascular disease pathophysiology and implications for treatment. *Prog Cardiovasc Dis*. 2003; 46:91-111.
- Bhujade R, Ibrahim T, Wanjpe AK, Chouhan DS. A comparative study to assess general health status and oral health score of tobacco users and nonusers in geriatric population in central India. *J Family Med Prim Care*. 2020;9(7):3387-3391. Published 2020 Jul 30. doi:10.4103/jfmpe.jfmpe\_157\_20.
- Bitzur R, Cohen H, Kamari Y, Shaish A, Harats D. Triglycerides and HDL cholesterol: stars or second leads in diabetes?. *Diabetes Care*. 2009;32 Suppl 2(Suppl 2):S373-S377. doi:10.2337/dc09-S343
- Bourassa KJ, Ruiz JM, Sbarra DA. Smoking and Physical Activity Explain the Increased Mortality Risk Following Marital Separation and Divorce: Evidence From the English Longitudinal Study of Ageing. *Ann Behav Med*. 2019;53(3):255-266. doi:10.1093/abm/kay038.
- Breenfeldt Andersen A, Bejder J, Bonne T, Olsen NV, Nordborg N. Repeated Wingate sprints is a feasible high-quality training strategy in moderate hypoxia. *PLoS One*. 2020;15(11):e0242439. Published 2020 Nov 13. doi:10.1371/journal.pone.0242439.
- Brunzell DH, Stafford AM, Dixon CI. Nicotinic receptor contributions to smoking: insights from human studies and animal models. *Curr Addict Rep*. 2015;2(1):33-46. doi:10.1007/s40429-015-0042-2.
- Buckler, Keith J; Turner, Philip J, Oxygen sensitivity of mitochondrial function in rat arterial chemoreceptor cells *PubMed Central*, 2013-01-01
- Bullen CH. Impact of tobacco smoking and smoking cessation on cardiovascular risk and disease. *Expert Rev Cardiovasc Ther*. 2008; 6(6):883-895.
- Buttar HS, Li T, Ravi N. Prevention of cardiovascular diseases: Role of exercise, dietary interventions, obesity and smoking cessation. *Exp Clin Cardiol*. 2005;10(4):229-249.
- Carbon Monoxide Specifically Inhibits Cytochrome C Oxidase of Human Mitochondrial Respiratory Chain September 2003 *Pharmacology & Toxicology* 93(3):142-6 DOI:10.1034/j.1600-0773.2003.930306.x SourcePubMed Authors: Jose Ramon Alonso Hospital Clínic de Barcelona.
- Centers for Disease Control and Prevention (US); National Center for Chronic Disease Prevention and Health Promotion (US); Office on Smoking and Health (US). Atlanta (GA): Centers for Disease Control and Prevention (US); 2010.
- Chu AJ. Tissue Factor, Blood Coagulation, and Beyond: An Overview. *Int J Inflam*. 2011; 2011:367284. doi: 10.4061/2011/367284. Epub 2011 Sep 20.

22. Csordas, A., Bernhard, D. The biology behind the atherothrombotic effects of cigarette smoke. *Nat Rev Cardiol* **10**, 219–230 (2013). <https://doi.org/10.1038/nrcardio.2013.8>
23. De Borja, A.T., Jost, R.T., Gass, R. et al. The influence of active and passive smoking on the cardiorespiratory fitness of adults. *Multidiscip Respir Med* **9**, 34 (2014). <https://doi.org/10.1186/2049-6958-9-34>
24. Dietrich T, Garcia RI, de Pablo P, Schulze PC, Hoffmann K. The effects of cigarette smoking on C-reactive protein concentrations in men and women and its modification by exogenous oral hormones in women. *Eur J Cardiovasc Prev Rehabil.* 2007; 14(5):694-700.
25. Dimitriadis G, Mitrou P, Lambadiari V, Maratou E, Raptis SA. Insulin effects in muscle and adipose tissue. *Diabetes Res Clin Pract.* 2011 Aug;93 Suppl 1:S52-9. doi: 10.1016/S0168-8227(11)70014-6. PMID: 21864752.
26. Duncker DJ, Bache RJ. Regulation of coronary blood flow during exercise. *Physiol Rev.* 2008 Jul;88(3):1009-86. doi: 10.1152/physrev.00045.2006. PMID: 18626066
27. Flammer AJ, Anderson T, Celermajer DS, et al. The assessment of endothelial function: from research into clinical practice. *Circulation.* 2012;126(6):753-767. doi:10.1161/CIRCULATIONAHA.112.093245.
28. G. Silveri, L. Pascasio, A. Miladinović, M. Ajčević and A. Accardo, Smoking effect on the circadian rhythm of blood pressure in hypertensive subjects, Department of Engineering and Architecture, University of Trieste, Trieste 34127, Italy 2 Department of Medical, Surgical and Health Care, CS of Geriatrics, University of Trieste, Trieste 34127, Italy GNB2020, June 10th-12th 2020
29. Gök, I; Celebi, I; Hacıoğlu, N; Ozic, C, Roles of beta2-adrenergic receptor gene polymorphisms in a Turkish population with obstructive sleep apnea syndrome or obesity. *Science.gov (United States)* 2014-10-20
30. Garcia M, Mulvagh SL, Merz CN, Buring JE, Manson JE. Cardiovascular Disease in Women: Clinical Perspectives. *Circ Res.* 2016;118(8):1273-1293. doi:10.1161/CIRCRESAHA.116.307547
31. George Papatheasiou, Anastasia Mamali, Spyridon Papafloratos and Efthimia Zerv Effects of smoking on cardiovascular function: the role of nicotine and carbon monoxide George Papatheasiou Prousis 22, Athens 17123 Greece
32. George, Effects of Smoking on Heart Rate at Rest and During Exercise, and on Heart Rate Recovery, in Young Adults may 2013 Hellenic journal of cardiology: HJC = Hellenikē kardiologikē epitheōrēsē 54(3):168-177.
33. Gepner AD, Piper ME, Johnson HM, Fiore MC, Baker TB, Stein JH. Effects of smoking and smoking cessation on lipids and lipoproteins: outcomes from a randomized clinical trial. *Am Heart J.* 2011;161(1):145-151.
34. Gille T, Sesé L, Aubourg E, et al. The Affinity of Hemoglobin for Oxygen Is Not Altered During COVID-19. *Front Physiol.* 2021;12:578708. Published 2021 Apr 12. doi:10.3389/fphys.2021.578708
35. Gusarov I, Shatalin K, Starodubtseva M, Nudler E. Endogenous nitric oxide protects bacteria against a wide spectrum of antibiotics. *Science.* 2009; 325:1380-1384. Harvard School of Public Health. The Greek Tobacco Epidemic. Centre for Global Tobacco Control. Boston, December 2011; available at: [www.smokefreegreece.org](http://www.smokefreegreece.org) (last day accessed 27 March 2013).
36. Harris KK, Zopey M, Friedman TC. Metabolic effects of smoking cessation [published correction appears in *Nat Rev Endocrinol.* 2016 Nov;12(11):684]. *Nat Rev Endocrinol.* 2016;12(5):299-308. doi:10.1038/nrendo.2016.32
37. Hogan MC, Bebout DE, Wagner PD. Effect of increased Hb-O<sub>2</sub> affinity on VO<sub>2</sub>max at constant O<sub>2</sub> delivery in dog muscle in situ. *J Appl Physiol* (1985). 1991 Jun;70(6):2656-62. doi: 10.1152/jappl.1991.70.6.2656. PMID: 1885462. 163(1):81-87.e1.
38. <https://aklectures.com/lecture/respiratory-system/carbon-monoxide-and-hemoglobin>
39. Hunter KA, Garlick PJ, Broom I, Anderson SE, McNurlan MA. Effects of smoking and abstention from smoking on fibrinogen synthesis in humans. *Clin Sci (Lond).* 2001; 100(4):459-465.
40. Komatsu, H., Yagasaki, K. & Yoshimura, K. Current nursing practice for patients on oral chemotherapy: a multicenter survey in Japan. *BMC Res Notes* **7**, 259 (2014). <https://doi.org/10.1186/1756-0500-7-259> (last day accessed 23 April 2014).
41. Kurt Brassington, Stanley Chan, Huei Seow, Aleksandar Dobric, Steven Bozinovski, Stavros Selemidis, and Ross Vlahos, Ebselen reduces cigarette smoke-induced vascular endothelial dysfunction in mice, IRMIT University, September 11, 2020.
42. Leaderer, B.P., Stolwijk, J.A.J. & Zgraniski, R.T. Health benefits due to reductions of CO levels. *Environmental Management* **1**, 131–137 (1977). <https://doi.org/10.1007/BF01866103>
43. Lande RG. Nicotine Addiction. Pathophysiology. Walter Reed Army Medical Center. Department of Psychiatry. Medscape Updated, December 13, 2012; available at: <http://emedicine.medscape.com/article/287555-overview#a0104> (last day accessed 27 October 2013).
44. Levine PH. An acute effect of cigarette smoking on platelet function. A possible link between smoking and arterial thrombosis. *Circulation.* 1973; 48(3):619-623.
45. Lusis AJ. Atherosclerosis. *Nature.* 2000;407(6801):233-241. doi:10.1038/3502520331. Mjos OD. Effect of free fatty acids on myocardial function and oxygen consumption in intact dogs. *J Clin Invest.* 1971; 50:1386- 1389.
46. Ma, B., Chen, Y., Wang, X. et al. Cigarette smoke exposure impairs lipid metabolism by decreasing low-density lipoprotein receptor expression in hepatocytes. *Lipids Health Dis* **19**, 88 (2020).
47. McGill HC. The cardiovascular pathology of smoking. *Am Heart J.* 1988; 115:250-257
48. Michael Pittilo R. Cigarette smoking, endothelial injury and cardiovascular disease. *Int J Exp Pathol.* 2000;81(4):219-230. doi:10.1046/j.1365-2613.2000.00162.x.
49. Miura H, Araki H, Matoba H, Kitagawa K. Relationship among oxygenation, myoelectric activity, and lactic acid accumulation in vastus lateralis muscle during exercise with constant work rate. *Int J Sports Med.* 2000 Apr;21(3):180-4. doi: 10.1055/s-2000-301. PMID: 10834349.
50. Myrna B. Schnur, MSN, RN, Systemic Vascular Resistance and Pulmonary Vascular Resistance: What's the Difference, May 25 2017
51. Papatheasiou G, Papandreou M, Galanos A, Kortianou E, Tsepis H, Kalfakakou V, et al. Smoking and physical activity interrelations in health science students. Is smoking associated with physical inactivity in young adults? *Hellenic J Cardiol.* 2012; 53:17-25.
52. Park SY, Gifford JR, Andbacka RH, et al. Cardiac, skeletal, and smooth muscle mitochondrial respiration: are all mitochondria created equal?. *Am J Physiol Heart Circ Physiol.* 2014;307(3):H346-H352. doi:10.1152/ajpheart.00227.2014.
53. Peto R, Lopez AD, Boreham J, Thun M. Mortality from smoking in developed countries 1950 -2000, 2nd edition: revised June 2006; available at: <http://www.ctsuo.ox.ac.uk/~tobacco/C0002.pdf> (last day accessed 27 March 2013).
54. Rafieian-Kopaei M, Setorki M, Doudi M, Baradaran A, Nasri H. Atherosclerosis: process, indicators, risk factors and new hopes. *Int J Prev Med.* 2014;5(8):927-946.

55. Roberts CK, Hevener AL, Barnard RJ. Metabolic syndrome and insulin resistance: underlying causes and modification by exercise training. *Compr Physiol*. 2013;3(1):1-58. doi:10.1002/cphy.c110062
56. Ross R, Glomset J, Harker L. Response to injury and atherogenesis. *Am J Pathol*. 1977; 86(3):675-84.
57. Sambola A, Osende J, Hathcock J, Degen M, Nemerson Y, Fuster V, et al. Role of Risk Factors in the Modulation of Tissue Factor Activity and Blood Thrombogenicity. *Circulation*. 2003; 107:973-977.
58. Sandoo A, van Zanten JJ, Metsios GS, Carroll D, Kitis GD. The endothelium and its role in regulating vascular tone. *Open Cardiovasc Med J*. 2010;4:302-312. Published 2010 Dec 23. doi:10.2174/1874192401004010302.
59. Tayade MC, Kulkarni NB. The effect of smoking on the cardiovascular autonomic functions: a cross sectional study. *J Clin Diagn Res*. 2013;7(7):1307-1310. doi:10.7860/JCDR/2013/5526.3133
60. U.S. Department of Health and Human Services. Public Health Service. How Tobacco Smoke Causes Disease: The Biology and Behavioural Basis for Smoking-Attributable Disease. A Report of the Surgeon General. USA, July 2013.
61. U.S. Department of Health and Human Services. National Centre for Chronic Disease Prevention and Health Promotion. Office on Smoking and Health. The Health Consequences of Smoking: A Report of the Surgeon General. Atlanta, 2004; available at: [http://www.cdc.gov/tobacco/data\\_statistics/sgr/2004/complete\\_report/index.htm](http://www.cdc.gov/tobacco/data_statistics/sgr/2004/complete_report/index.htm) (last day accessed 22 September 2013)
62. USA Institute of Medicine of the National Academies. Secondhand Smoke Exposure and Cardiovascular Effects: Making Sense of the Evidence. Washington DC: The National Academies Press, National Academy of Sciences, July 2013.
63. Yun AJ, Bazar KA, Lee PY, Gerber A, Daniel SM. The smoking gun: many conditions associated with tobacco exposure may be attributable to paradoxical compensatory autonomic responses to nicotine. *Med Hypotheses*. 2005;64(6):1073-9. doi: 10.1016/j.mehy.2004.11.040. PMID: 15823687.
64. Wan Ahmad WN, Sakri F, Mokhsin A, Rahman T, Mohd Nasir N, Abdul-Razak S, Md Yasin M, Mohd Ismail A, Ismail Z, Nawawi H. Low serum high density lipoprotein cholesterol concentration is an independent predictor for enhanced inflammation and endothelial activation. *PLoS One*. 2015 Jan 23;10(1):e0116867.
65. World Health Organization. Report on the Global Tobacco Epidemic. Geneva, 2008; available at: <http://www.who.int/tobacco/mpower/mpow> (October 2013).
66. World Health Organization. Regional Office for Europe. The European Tobacco Control Report 2007. Geneva, October 2013.
67. World Health Organization. World Health Statistics. Geneva, 2011; available at: [http://www.who.int/gho/publications/world\\_health\\_statistics/2011/en/](http://www.who.int/gho/publications/world_health_statistics/2011/en/) (last day accessed 24 September 2013).



# ЖУРНАЛ КАРДИОРЕСПИРАТОРНЫХ ИССЛЕДОВАНИЙ

## JOURNAL OF CARDIORESPIRATORY RESEARCH

**№2 (2021)**

**Контакт редакций журналов. [www.tadqiqot.uz](http://www.tadqiqot.uz)**  
ООО Тадqiqот город Ташкент,  
улица Амира Темура пр.1, дом-2.  
Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Тел: (+998-94) 404-0000

**Editorial staff of the journals of [www.tadqiqot.uz](http://www.tadqiqot.uz)**  
Tadqiqot LLC the city of Tashkent,  
Amir Temur Street pr.1, House 2.  
Web: <http://www.tadqiqot.uz/>; Email: [info@tadqiqot.uz](mailto:info@tadqiqot.uz)  
Phone: (+998-94) 404-0000